

# ABDOMINAL PAIN

- Location
- Work-up
- Acute pain syndromes
- Chronic pain syndromes

# Epigastric Pain

- PUD
- GERD
- MI
- AAA- abdominal aortic aneurysm
- Pancreatic pain
- Gallbladder and common bile duct obstruction

# Right Upper Quadrant Pain

- Acute Cholecystitis and Biliary Colic
- Acute Hepatitis or Abscess
- Hepatomegaly due to CHF
- Perforated Duodenal Ulcer
- Herpes Zoster
- Myocardial Ischemia
- Right Lower Lobe Pneumonia



# Left Upper Quadrant Pain

- Acute Pancreatitis
- Gastric ulcer
- Gastritis
- Splenic enlargement, rupture or infarction
- Myocardial ischemia
- Left lower lobe pneumonia



# Right lower Quadrant Pain

- Appendicitis
- Regional Enteritis
- Small bowel obstruction
- Leaking Aneurysm
- Ruptured Ectopic Pregnancy
- PID
- Twisted Ovarian Cyst
- Ureteral Calculi
- Hernia

# Left Lower Quadrant Pain

- Diverticulitis
- Leaking Aneurysm
- Ruptured Ectopic pregnancy
- PID
- Twisted Ovarian Cyst
- Ureteral Calculi
- Hernia
- Regional Enteritis

# Periumbilical Pain

- Disease of transverse colon
- Gastroenteritis
- Small bowel pain
- Appendicitis
- Early bowel obstruction

# Diffuse Pain

- Generalized peritonitis
- Acute Pancreatitis
- Sickle Cell Crisis
- Mesenteric Thrombosis
- Gastroenteritis
- Metabolic disturbances
- Dissecting or Rupturing Aneurysm
- Intestinal Obstruction
- Psychogenic illness





# Referred Pain

- Pneumonia (lower lobes)
- Inferior myocardial infarction
- Pulmonary infarction



# TYPES OF ABDOMINAL PAIN

- Visceral
  - originates in abdominal organs covered by peritoneum
- Colic
  - crampy pain
- Parietal
  - from irritation of parietal peritoneum
- Referred
  - produced by pathology in one location felt at another location

# ORGANIC VERSUS FUNCTIONAL PAIN

| <u>HISTORY</u> | <u>ORGANIC</u> | <u>FUNCTIONAL</u> |
|----------------|----------------|-------------------|
|----------------|----------------|-------------------|

|                |  |                       |
|----------------|--|-----------------------|
| Pain character | Acute, persistent pain increasing in intensity | Less likely to change |
|----------------|--|-----------------------|

|                   |                   |                   |
|-------------------|-------------------|-------------------|
| Pain localization | Sharply localized | Various locations |
|-------------------|-------------------|-------------------|

|                           |                  |           |
|---------------------------|------------------|-----------|
| Pain in relation to sleep | Awakens at night | No affect |
|---------------------------|------------------|-----------|

|                               |              |              |
|-------------------------------|--------------|--------------|
| Pain in relation to umbilicus | Further away | At umbilicus |
|-------------------------------|--------------|--------------|

|                     |  |   |
|---------------------|--|---|
| Associated symptoms | Fever, anorexia, vomiting, wt loss, anemia, elevated ESR | Headache, dizziness, multiple system complaints |
|---------------------|--|---|

|                      |               |         |
|----------------------|---------------|---------|
| Psychological stress | None reported | Present |
|----------------------|---------------|---------|

⋮

# WORK-UP OF ABDOMINAL PAIN

## HISTORY

- Onset
- Qualitative description
- Intensity
- Frequency
- Location - Does it go anywhere (referred)?
- Duration
- Aggravating and relieving factors



# WORK-UP

## PHYSICAL EXAMINATION

- Inspection
- Auscultation
- Percussion
- Palpation
- Guarding - rebound tenderness
- Rectal exam
- Pelvic exam



•  
•  
•

# WORK-UP

## LABORATORY TESTS

- U/A
- CBC
- Additional depending on rule outs
  - amylase, lipase, LFT's



# WORK-UP

## DIAGNOSTIC STUDIES

- Plain X-rays (flat plate)
- Contrast studies - barium (upper and lower GI series)
- Ultrasound
- CT scanning
- Endoscopy
- Sigmoidoscopy, colonoscopy



# Common Acute Pain Syndromes

- Appendicitis
- Acute diverticulitis
- Cholecystitis
- Pancreatitis
- Perforation of an ulcer
- Intestinal obstruction
- Ruptured AAA
- Pelvic disorders



# APPENDICITIS

- Inflammatory disease of wall of appendix
- Diagnosis based on history and physical
- Classic sequence of symptoms
  - abdominal pain (begins epigastrium or periumbilical area, anorexia, nausea or vomiting)
  - followed by pain over appendix and low grade fever

# DIAGNOSIS

- Physical examination
  - low grade fever
  - McBurney's point
  - rebound, guarding, +psoas sign
- CBC, HCG
  - WBC range from 10,000-16,000

SURGERY

# DIVERTICULITIS

- Results from stagnation of fecal material in single diverticulum leading to pressure necrosis of mucosa and inflammation
- Clinical presentation
  - most pts have h/o diverticula
  - mild to moderate, colicky to steady, aching abdominal pain - usually LLQ
  - may have fever and leukocytosis



# CHOLECYSTITIS

- Results from obstruction of cystic or common bile duct by large gallstones
- Colicky pain with progression to constant pain in RUQ that may radiate to R scapula
- Physical findings
  - tender to palpation or percussion RUQ
  - may have palpable gallbladder

•  
•  
•

## DIAGNOSIS

- CBC, LFTs (bilirubin, alkaline phosphatase), serum pancreatic enzymes
- Plain abdominal films demonstrate biliary air hepatomegaly, and maybe gallstones
- Ultrasound - considered accurate about 95%

## MANAGEMENT

- Admission

# PANCREATITIS

- History of cholelithiasis or ETOH abuse
- Pain steady and boring, unrelieved by position change - LUQ with radiation to back - nausea and vomiting, diaphoretic
- Physical findings;
  - acutely ill with abdominal distention, ↓ BS
  - diffuse rebound
  - upper abd may show muscle rigidity

- 
- 
- 

## • Diagnostic studies

- CBC

- Ultrasound

- Serum amylase and lipase

- amylase rises 2-12 hours after onset and returns to normal in 2-3 days

- lipase is elevated several days after attack

## Management

- Admission



# PEPTIC ULCER PERFORATION

- Life-threatening complication of peptic ulcer disease - more common with duodenal than gastric
- Predisposing factors
  - *Helicobacter pylori* infections
  - NSAIDs
  - hypersecretory states

- Sudden onset of severe intense, steady epigastric pain with radiation to sides, back, or right shoulder
- Past h/o burning, gnawing pain worse with empty stomach
- Physical findings
  - epigastric tenderness
  - rebound tenderness
  - abdominal muscle rigidity
- Diagnostic studies
  - upright or lateral decubitus X-ray shows air under the diaphragm or peritoneal cavity

**REFER - SURGICAL EMERGENCY**

# SMALL BOWEL OBSTRUCTION

- Distention results in decreased absorption and increased secretions leading to further distention and fluid and electrolyte imbalance
- Number of causes
- Sudden onset of crampy pain usually in umbilical area of epigastrium - vomiting occurs early with small bowel and late with large bowel

- Physical findings

- hyperactive, high-pitched BS
- fecal mass may be palpable
- abdominal distention
- empty rectum on digital exam

- Diagnosis

- CBC
- serum amylase
- stool for occult blood
- type and crossmatch
- abdominal X-ray

- Management

- Hospitalization

# RUPTURED AORTIC ANEURYSM

- AAA is abnormal dilation of abdominal aorta forming aneurysm that may rupture and cause exsanguination into peritoneum
- More frequent in elderly
- Sudden onset of excruciating pain may be felt in chest or abdomen and may radiate to legs and back

- 
- 
- 
- 
- **Physical findings**

- appears shocky
- VS reflect impending shock
- deficit or difference in femoral pulses

- **Diagnosis**

- CT or MRI
- ECG, cardiac enzymes

**SURGICAL EMERGENCY**

•  
•  
•

# PELVIC PAIN

- Ectopic pregnancy
- PID
- UTI
- Ovarian cysts

# CHRONIC PAIN SYNDROMES

- Irritable bowel syndrome
- Chronic pancreatitis
- Diverticulosis
- Gastroesophageal reflux disease (GERD)
- Inflammatory bowel disease
- Duodenal ulcer
- Gastric ulcer



•  
•  
•

# IRRITABLE BOWEL SYNDROME

- GI condition classified as functional as no identifiable structural or biochemical abnormalities
- Affects 14%-24% of females and 5%-19% of males
- Onset in late adolescence to early adulthood
- Rare to see onset > 50 yrs old

# SYMPTOMS

- Pain described as nonradiating, intermittent, crampy located lower abdomen
- Usually worse 1-2 hrs after meals
- Exacerbated by stress
- Relieved by BM
- Does not interrupt sleep
  - critical to diagnosis of IBS

# DIAGNOSIS

## ROME DIAGNOSTIC CRITERIA

- **3 month minimum of following symptoms in continuous or recurrent pattern**

Abdominal pain or discomfort relieved by BM & associated with either:

Change in frequency of stools

*and/or*

Change in consistency of stools

•  
•  
•

**Two or more of following symptoms on  
25% of occasions/days:**

Altered stool frequency

>3 BMs daily or <3 BMs/week

Altered stool form

Lumpy/hard or loose/watery

Altered stool passage

Straining, urgency, or feeling of incomplete  
evacuation

Passage of mucus

Feeling of bloating or abdominal distention

# DIAGNOSTIC TESTS

- Limited - R/O organic disease
- CBC with diff
- ESR
- Electrolytes
- BUN, creatinine
- TSH
- Stool for occult blood and O & P
- Flexible sigmoidoscopy

# MANAGEMENT

- Goals of management
  - exclude presence of underlying organic disease
  - provide support, support, & reassurance
- Dietary modification
- Pharmacotherapy
- Alternative therapies

•  
•  
•

Physician consultation is indicated if initial treatment of IBS fails, if organic disease is suspected, and/or if the patient who presents with a change in bowel habits is over 50

# CHRONIC PANCREATITIS

- Alcohol major cause
- Malnutrition - outside US
- Patients >40 yrs with pancreatic dysfunction must be evaluated for pancreatic cancer
- Dysfunction between 20 to 40 yrs old R/O cystic fibrosis
- 50% of pts with chronic pancreatitis die within 25 yrs of diagnosis



# SYMPTOMS

- Pain - may be absent or severe, recurrent or constant
- Usually abdominal, sometimes referred upper back, anterior chest, flank
- Wt loss, diarrhea, oily stools
- N, V, or abdominal distention less reported



# DIAGNOSIS

- CBC
- Serum amylase (present during acute attacks)
- Serum lipase
- Serum bilirubin
- Serum glucose
- Serum alkaline phosphatase
- Stool for fecal fat
- CT scan



# MANAGEMENT

- Should be comanaged with a specialist
- Pancreatic dysfunction
  - diabetes
  - steatorrhea & diarrhea
  - enzyme replacement

# DIVERTICULOSIS

- Uncomplicated disease, either asymptomatic or symptomatic
- Considered a deficiency disease of 20th century Western civilization
- Rare in first 4 decades - occurs in later years
- Incidence - 50% to 65% by 80 years

# SYMPTOMS

- 80% - 85% remain symptomless - found by diagnostic study for other reason
- Irregular defecation, intermittent abdominal pain, bloating, or excessive flatulence
- Change in stool - flattened or ribbonlike
- Recurrent bouts of steady or crampy pain
- May mimic IBS except older age

•  
•  
•

# DIAGNOSIS

- CBC
- Stool for occult blood
- Barium enema

# MANAGEMENT

- Increased fiber intake - 35 g/day
- Increase fiber intake gradually
- Avoid
  - popcorn
  - corn
  - nuts
  - seeds

# GASTROESOPHAGEAL REFLUX DISEASE

- Movement of gastric contents from stomach to esophagus
- May produce S & S within esophagus, pharynx, larynx, respiratory tract
- Most prevalent condition affecting GI tract
- About 15% of adults use antacid > 1x/wk



# SYMPTOMS

- Heartburn - most common (severity of does not correlate with extent of tissue damage)
- Burning, gnawing in mid-epigastrium worsens with recumbency
- Water brash (appearance of salty-tasting fluid in mouth because stimulate saliva secretion)
- Occurs after eating may be relieved with antacids (occurs within 1 hr of eating - usually large meal of day)

- Dysphagia & odynophagia predictive of severe disease
- Chest pain - may mimic angina
- Foods that may precipitate heartburn
  - high fat or sugar
  - chocolate, coffee, & onions
  - citrus, tomato-based, spicy
- Cigarette smoking and alcohol
- Aspirin, NSAIDS, potassium, pills

# DIAGNOSIS

- History of heartburn without other symptoms of serious disease
- Empiric trial of medication without testing
- Testing for those who do have persistent or unresponsive heartburn or signs of tissue injury
- CBC, *H. pylori* antibody
- Barium swallow
- Endoscopy for severe or atypical symptoms

# MANAGEMENT

- Lifestyle changes
  - smoking cessation
  - reduce ETOH consumption
  - reduce dietary fat
  - decreased meal size
  - weight reduction
  - elevate head of bed 6 inches

- - 
  -
- elimination of medications that are mucosal irritants or that lower esophageal pressure
  - avoidance of chocolate, peppermint, coffee, tea, cola beverages, tomato juice, citrus fruit juices
  - avoidance of supine position for 2 hours after meal
  - avoidance of tight fitting clothes

# MEDICATIONS

- Antacids with lifestyle changes may be sufficient
- H<sub>2</sub>-histamine receptor antagonists in divided doses
  - approximately 48% of pts with esophagitis will heal on this regimen
  - tid dosing more effective for symptom relief and healing
  - long-term use is appropriate



- Proton pump inhibitors - prilosec & prevacid
  - once a day dosing
  - compared with H<sub>2</sub>RA have greater efficacy relieving symptoms & healing
  - treat moderate to severe for 8 wks
  - may continue with maintenance to prevent relapse



# MAINTENANCE THERAPY

- High relapse rate - 50% within 2 months, 82% within 6 months without maintenance
- If symptoms return after treatment need maintenance
- Full dose H<sub>2</sub>RA for most patients with nonerosive GERD
- Proton pump inhibitors for severe or complicated



# INFLAMMATORY BOWEL DISEASE

- Chronic inflammatory condition involving intestinal tract with periods of remission and exacerbation
- Two types
  - Ulcerative colitis (UC)
  - Crohn's disease

# ULCERATIVE COLITIS

- Chronic inflammation of colonic mucosa
- Inflammation diffuse & continuous beginning in rectum
- May involve entire colon or only rectum (proctitis)
- Inflammation is continuous

# CROHN'S DISEASE

- Chronic inflammation of all layers on intestinal tract
- Can involve any portion from mouth to anus
- 30%-40% small intestine (ileitis)
- 40%-45% small & large intestine (ileocolitis)
- 15%-25% colon (Crohn's colitis)
- Inflammation can be patchy

- Annual incidence of UC & Crohn's similar in both age of onset & worldwide distribution
- About 20% more men have UC
- About 20% more women have Crohn's
- Peak age of onset - between 15 & 25 yrs

# SYMPTOMS

- Both have similar presentations
- Abdominal pain may be only complaint and may have been intermittent for years
- Abdominal pain and diarrhea present in most pts
- Pain diffuse or localized to RLQ-LLQ
- Cramping sensation - intermittent or constant

- Tenesmus & fecal incontinence
- Stools loose and/or watery - may have blood
- Rectal bleeding common with colitis
- Other complaints
  - fatigue
  - weight loss
  - anorexia
  - fever, chills
  - nausea, vomiting
  - joint pains
  - mouth sores

# PHYSICAL EXAMINATION

- May be in no distress to acutely ill
- Oral aphthous ulcers
- Tender lower abdomen
- Hyperactive bowel sounds
- Stool for occult blood may be +
- Perianal lesions
- Need to look for fistulas & abscesses

# DIAGNOSIS

- CBC
- Stool for culture, ova & parasites, *C. difficile*
- Stool for occult blood
- Flexible sigmoidoscopy - useful to determine source of bright red blood
- Colonoscopy with biopsy
- Endoscopy may show “skip” areas
- May be difficult to distinguish one from other



# MANAGEMENT

- Should be comanaged with GI
- 5-aminosalicylic acid products
- Corticosteroids
- Immunosuppressives
- Surgery



# DUODENAL ULCERS

- Incidence increasing secondary to increasing use of NSAIDs, *H. pylori* infections
- Imbalance both in amount of acid-pepsin production delivered from stomach to duodenum and ability of lining to protect self





# RISK FACTORS

- Stress
- Cigarette smoking
- COPD
- Alcohol
- Chronic ASA & NSAID use



# GENETIC FACTORS

- Zollinger-Ellison syndrome
- First degree relatives with disease
- Blood group O
- Elevated levels of pepsinogen I
- Presence of HLA-B5 antigen
- Decreased RBC acetylcholinesterase

# INCIDENCE

- About 16 million individuals will have during lifetime
- More common than gastric ulcers
- Peak incidence; 5th decade for men, 6th decade for women
- 75%-80% recurrence rate within 1yr of diagnosis without maintenance therapy
- >90% of duodenal ulcers caused by *H.pylori*

# SYMPTOMS

- Epigastric pain
- Sharp, burning, aching, gnawing pain occurring 1 □ - 3 hrs after meals or in middle of night
- Pain relieved with antacids or food
- Symptoms recurrent lasting few days to months
- Weight gain not uncommon

# DIAGNOSIS

- CBC
- Serum for *H. pylori*
- Stool for occult blood

# MANAGEMENT

- 2 week trial of antiulcer med - d/c NSAIDs
- If *H. pylori* present - treat
- If no *H. pylori* & symptoms do not resolve after 2 wks refer to GI for endoscopy
- Antiulcer meds
  - H<sub>2</sub>RA; associated with 75%-90% healing over 4-6week period followed by 1 yr maintenance
  - inhibits P-450 pathway; drug interactions



# MANAGEMENT (CONT)

- Proton pump inhibitors
  - daily dosing
  - documented improved efficacy over H<sub>2</sub>-RA blockers
- Prostaglandin therapy - misoprostol
  - use with individuals who cannot d/c NSAIDs

# GASTRIC ULCERS

- *H. pylori* identified in 65% to 75% of patients with non-NSAID use
- 5% - 25% of patients taking ASA/NSAID develop gastric ulcers (inhibits synthesis of prostaglandin which is critical for mucosal defense)
- Malignancy cause of

# OTHER RISK FACTORS

- Caffeine/coffee
- Alcohol
- Smoking
- First-degree relative with gastric ulcer

# SYMPTOMS

- Pain similar to duodenal but may be increased by food
- Location - LUQ radiating to back
- Bloating, belching, nausea, vomiting, weight loss
- NSAID-induced ulcers usually painless - discovered secondary to melena or iron deficiency anemia

•  
•  
•

# DIAGNOSIS

- CBC
- Serum for *H. pylori*
- Carbon-labeled breath test
- Stool for occult blood
- Endoscopy

# MANAGEMENT

- Treat *H.pylori* if present
- Proton pump inhibitors shown to be superior to H<sub>2</sub>-RA
- Need to use proton pump inhibitor for up to 8 wks
- Do not need maintenance if infection eradicated and NSAIDs d/c'd
- Consider misoprostol if cannot d/c NSAID