



Maternal health

Hafsa Raheel

Associate Professor & Consultant

Preventive medicine and Public health

Department of Family and Community Medicine

KSU

Objectives

- Understand the maternal health issues globally
- Understand the causes of maternal deaths and mortality
- Understand the interventions done globally to decrease maternal deaths and morbidity
 - Antenatal care
 - Promotion of breast feeding practices.....BFHI
- Discuss and understand what preventive services for maternal health are delivered in KSA

MATERNAL HEALTH




Defination

Maternal health refers to the health of women during pregnancy, childbirth and the postpartum period. While motherhood is often a positive and fulfilling experience, for too many women it is associated with suffering, ill-health and even death.

Fast Facts about Maternal Health . . . WHO Fact sheet (2021)

- **Every day in 2017, approximately 810 women died from preventable causes related to pregnancy and childbirth.**
- **Between 2000 and 2017, the maternal mortality ratio (MMR, number of maternal deaths per 100,000 live births) dropped by about 38% worldwide.**
- **94% of all maternal deaths occur in low and lower middle-income countries.**
- **Young adolescents (ages 10-14) face a higher risk of complications and death as a result of pregnancy than other women.**
- **Skilled care before, during and after childbirth can save the lives of women and new-borns.**

Maternal mortality ratio (per 100 000 live births), 2015



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.


Data Source: World Health Organization
 Map Production: Health Statistics and Information Systems (HSI)
 World Health Organization



© WHO 2015. All rights reserved.


Maternal death: the death of a woman while pregnant, (or within 42 days of termination of pregnancy).

- irrespective of the duration and the site of the pregnancy




Maternal death: the death of a woman while pregnant, (or within 42 days of termination of pregnancy).

- All maternal deaths are either direct obstetric deaths or indirect obstetric deaths



Maternal death: the death of a woman while pregnant, (or within 42 days of termination of pregnancy).

Accidental or incidental causes of death are not classified as maternal deaths.






Why women are dying?

Women die as a result of complications during and following pregnancy and childbirth.

- The major complications that account for nearly 75% of all maternal deaths are:
 - severe bleeding (mostly bleeding after childbirth)
 - infections (usually after childbirth)
 - high blood pressure during pregnancy (pre-eclampsia and eclampsia)
 - complications from delivery
 - unsafe abortion
- The remainder are caused by or associated with diseases such as malaria, and AIDS during pregnancy.

Global Causes of Maternal Mortality



Ref: Say L et al., 'Global causes of maternal death: a WHO systematic analysis' Lancet Global Health. [http://dx.doi.org/10.1016/S2214-109X\(14\)70227-X](http://dx.doi.org/10.1016/S2214-109X(14)70227-X), May 6, 2014.

A 3D grid of spheres on a blue background. The spheres are arranged in a regular, repeating pattern that recedes into the distance, creating a sense of depth. The background is a solid, dark blue color.

Why do women not get the care they need?

Why do these women die?

Three Delays Model


- Delay in decision to seek care
 - Lack of understanding of complications
 - Acceptance of maternal death
 - Low status of women
 - Socio-cultural barriers to seeking care
- Delay in reaching care
 - Mountains, islands, rivers — poor organization
- Delay in receiving care
 - Supplies, personnel
 - Poorly trained personnel with punitive attitude
 - Finances

Trends in maternal mortality 1990 - 2015


Maternal mortality fell by almost half between 1990 and 2015

Maternal mortality ratio (maternal deaths per 100,000 live births in women aged 15 to 49), by region, 1990, 2010 and 2015

Ref:
<http://data.unicef.org/maternal-health/maternal-mortality.html#sthash.Eu3mJpN1.dpuf>



Maternal mortality ratio
 (maternal deaths per 100 000 live births)
 Globally and by WHO region, 1990–2015



Where do Maternal Mortality data come from?


- Vital registration data - **MM Rate and MM Ratio**
- Health service data – maternity registers - **MM Ratio**
- Special studies
 - Hospital studies – tracing deaths, interviews
 - Research, longitudinal studies, verbal autopsy
- Surveys & censuses
 - Direct estimation - **Rate and Ratio**
 - Sisterhood method (indirect) – **Rate and Ratio**

Maternal Mortality Indicators

- Maternal mortality ratio
- Maternal mortality rate
- Life-time risk of maternal mortality
- Proportion maternal

Maternal mortality ratio:
the number of maternal
deaths per *live births*

Numerator: Maternal deaths



Denominator: Live births




Maternal mortality rate:

the number of maternal deaths in a given period per population of *women who are of reproductive age*

Numerator: Maternal deaths

Denominator: Women of reproductive age



Other Maternal Mortality Indicators

- **Life time risk of maternal mortality** = (N of maternal deaths over the reproductive life span) / (women entering the reproductive period)
- **Proportion maternal** = proportion of all female deaths due to maternal causes = (N of maternal deaths in a period/Number of all female deaths in same period) * 100

Why has the maternal mortality declined?

Global response ???

1 NO POVERTY



2 ZERO HUNGER



3 GOOD HEALTH AND WELL-BEING



4 QUALITY EDUCATION



5 GENDER EQUALITY



6 CLEAN WATER AND SANITATION



7 AFFORDABLE AND CLEAN ENERGY



8 DECENT WORK AND ECONOMIC GROWTH



9 INDUSTRY, INNOVATION AND INFRASTRUCTURE



10 REDUCED INEQUALITIES



11 SUSTAINABLE CITIES AND COMMUNITIES



THE GLOBAL GOALS

For Sustainable Development


12 RESPONSIBLE CONSUMPTION AND PRODUCTION



13 CLIMATE ACTION



14 LIFE BELOW WATER



15 LIFE ON LAND



16 PEACE AND JUSTICE STRONG INSTITUTIONS



17 PARTNERSHIPS FOR THE GOALS



Global response

● Sustainable Development Goal 3

- 3.1 By 2030, reduce the global maternal mortality ratio to less than 70 per 100 000 live births.

Successful Interventions for Maternal Care

Antenatal care

- Nutrition support (anemia, adequate caloric intake)
- Personal hygiene, dental care, rest (2 hrs) and sleep (8 hrs), regular bowel habits..enough fiber and fruit intake...avoid constipation
- Immunization (mother and the new born)
- Drugs; thalidomide (deformed hands), corticosteroids (impair fetal growth), streptomycin (8th nerve damage)
- Education on delivery and care of the new born
- Identifying high risk pregnancies, smoking and exposure to passive smoking

Antenatal care.....cont

- Emphasizing on ANC visits and maintenance of AN card
- Importance and management of lactation
(importance/benefits of breast feeding, exclusive breast feeding, problems arising from breast feeding)
- Advise on birth spacing

Ref: WHO recommendations on maternal health, guidelines to improve maternal health. 2017.

Available at:<http://apps.who.int/iris/bitstream/10665/259268/1/WHO-MCA-17.10-eng.pdf?ua=1>

Why is ANC critical?

Through timely and appropriate evidence-based actions related to health promotion, disease prevention, screening, and treatment

- Reduces complications from pregnancy and childbirth

- Reduces stillbirths and perinatal deaths

- Integrated care delivery throughout pregnancy

2016 WHO ANC model

WHO FANC model	2016 WHO ANC model
<i>First trimester</i>	
Visit 1: 8-12 weeks	Contact 1: up to 12 weeks
<i>Second trimester</i>	
Visit 2: 24-26 weeks	Contact 2: 20 weeks Contact 3: 26 weeks
<i>Third trimester</i>	
Visit 3: 32 weeks Visit 4: 36-38 weeks	Contact 4: 30 weeks Contact 5: 34 weeks Contact 6: 36 weeks Contact 7: 38 weeks Contact 8: 40 weeks
Return for delivery at 41 weeks if not given birth.	

Antenatal care

A 3D grid of spheres on a blue background. The spheres are arranged in a regular, repeating pattern that recedes into the distance, creating a sense of depth. The background is a solid, dark blue color.

History taking (1st visit)


- Confirm the pregnancy
- Any previous complications (abortions, still births)
- Calculate LMP (add 9 months and 7 days to the first day of menstruation)
- Record symptoms; fever, vomiting, abnormal vaginal bleeding, palpitation, easy fatigability, breathlessness, generalized swelling, burning micturition, decreased or absent fetal movements
- Any concurrent illness; asthma, heart disease, jaundice, HTN, DM, TB, HIV, STIs, thalassemia, bleeding disorders
- Family history of twins, congenital malformations
- History of drug allergies, or drugs

Physical exam

- **General physical**; pallor, pulse (N 60 – 90 mins), respiratory rate (N 18-20 breaths/min), edema (slight edema is normal, if co-existent with any diseases eg HTN, referral)
- BP (every visit)
 - High BP; ≥ 2 readings 140/90
 - Urine +2 albumin
 - High BP + albuminuria = pre-eclampsia ---refer
- Weight ; 9-11 kg during pregnancy. Approx. 2 kg /month
- Breast exam

Abdominal exam

● Fundal height



At about three months (13-14 weeks), the top of the uterus is usually just above the mother's pubic bone (where her pubic hair begins).

At about five months (20-22 weeks), the top of the uterus is usually right at the mother's bellybutton (umbilicus or navel).

At about eight to nine months (36-40 weeks), the top of the uterus is almost up to the bottom of the mother's ribs.

Babies may drop lower in the weeks just before birth. You can look back at Figure 7.1 in Study Session 7 to see a diagram of fundal height at various weeks of gestation.

Assessment of gestational age

- Routine US + LMP (history)

- **Lab investigations:**

- Pregnac test, Hb estimation, Urine for albumin and sugar, blood grouping, Rh factor, VDRL, HIV testing, Blood sugar, HBsAg for Hep B

Ultrasound

● Fetal assessment

- One ultrasound scan before 24 weeks of gestation (**early ultrasound**) is recommended for pregnant women to estimate gestational age
 - Advantages; improve detection of fetal anomalies and multiple pregnancies, reduce induction of labour for post-term pregnancy, and improve a woman's pregnancy experience.

Antenatal care counseling

● Nutritional recommendations:

- Counselling about healthy eating and keeping physically active during pregnancy is recommended for pregnant women to stay healthy and to prevent excessive weight gain during pregnancy.
- Daily oral iron and folic acid supplementation with **30 mg to 60 mg of elemental iron** and **400 µg (0.4 mg) of folic acid** is recommended for pregnant women to prevent maternal anaemia, puerperal sepsis, low birth weight, and preterm birth
- Foods rich in iron; dates, green leafy vegetables, red beans, gauvas, red meats

Antenatal care

● Maternal assessment

- Hyperglycaemia first detected at any time during pregnancy should be classified as either gestational diabetes mellitus (GDM) or diabetes mellitus in pregnancy.
- Health-care providers should ask all pregnant women about their tobacco use (past and present) and exposure to second-hand smoke as early as possible in the pregnancy and at every antenatal care visit.
- At every visit, history of TB, HIV, and alcohol intake should also be accessed....in high prevalence areas

Preventive services

- A seven-day antibiotic regimen is recommended for all pregnant women with **asymptomatic bacteriuria (ASB)** to prevent persistent bacteriuria, preterm birth and low birth weight
- **Tetanus toxoid vaccination** is recommended for all pregnant women, depending on previous tetanus vaccination exposure, to prevent neonatal mortality from tetanus.

Tetanus vaccination

Table 2 Guidelines for tetanus toxoid immunization of women who were immunized during infancy, childhood or adolescence^b

Age at last vaccination	Previous immunizations (based on written records)	Recommended Immunizations	
		At present contact/pregnancy	Later (at intervals of at least one year)
Infancy	3 DTP	2 doses of TT/Td (min.4 weeks interval between doses)	1 dose of TT/Td
Childhood	4 DTP	1 dose of TT/Td	1 dose of TT/Td
School age	3 DTP + 1 DT/Td	1 dose of TT/Td	1 dose of TT/Td
School age	4 DTP + 1 DT/Td	1 dose of TT/Td	None
Adolescence	4 DTP + 1 DT at 4-6 yrs + 1 TT/Td at 14-16 yrs	None	None

^b Adapted from: Galazka AM. *The immunological basis for immunization series. Module 3: tetanus*. Geneva, World Health Organization, 1993 (WHO/EPI/GEN/93.13), page 17.

Table 1 Tetanus toxoid immunization schedule for women of childbearing age and pregnant women without previous exposure to TT, Td or DTP^a

Dose of TT or Td (according to card or history)	When to give	Expected duration of protection
1	At first contact or as early as possible in pregnancy	None
2	At least 4 weeks after TT1	1-3 years
3	At least 6 months after TT2 or during subsequent pregnancy	At least 5 years
4	At least one year after TT3 or during subsequent pregnancy	At least 10 years
5	At least one year after TT4 or during subsequent pregnancy	For all childbearing age years and possibly longer

^a Source: *Core information for the development of immunization policy. 2002 update*. Geneva. World Health Organization, 2002 (document WHO/V&B/02.28), page 130.

Common physiological symptoms

Ginger, chamomile, vitamin B6 and/or acupuncture are recommended for the **relief of nausea** in early pregnancy, based on a woman's preferences and available options.

Advice on diet and lifestyle is recommended to prevent and relieve **heartburn** in pregnancy. Antacid preparations can be offered to women with troublesome symptoms that are not relieved by lifestyle modification.

Magnesium, calcium or non-pharmacological treatment options can be used for the relief of **leg cramps** in pregnancy, based on a woman's preferences and available options.

Regular exercise throughout pregnancy is recommended to prevent **low back and pelvic pain**. There are a number of different treatment options that can be used, such as physiotherapy, support belts and acupuncture, based on a woman's preferences and available options.

Wheat bran or other fibre supplements can be used to relieve **constipation** in pregnancy if the condition fails to respond to dietary modification, based on a woman's preferences and available options.

Non-pharmacological options, such as compression stockings, leg elevation and water immersion, can be used for the management of **varicose veins and oedema** in pregnancy, based on a woman's preferences and available options.

Baby friendly hospital initiative (BFHI)

- The Baby-friendly Hospital Initiative (BFHI) was launched by WHO and UNICEF in 1991.
- The initiative is a global effort to implement practices that protect, promote and support breastfeeding.

The TEN STEPS to Successful Breastfeeding

1 HOSPITAL POLICIES

Hospitals support mothers to breastfeed by...



2 STAFF COMPETENCY

Hospitals support mothers to breastfeed by...



3 ANTENATAL CARE

Hospitals support mothers to breastfeed by...



4 CARE RIGHT AFTER BIRTH

Hospitals support mothers to breastfeed by...



5 SUPPORT MOTHERS WITH BREASTFEEDING

Hospitals support mothers to breastfeed by...



6 SUPPLEMENTING

Hospitals support mothers to breastfeed by...



7 ROOMING-IN

Hospitals support mothers to breastfeed by...



8 RESPONSIVE FEEDING

Hospitals support mothers to breastfeed by...



9 BOTTLES, TEATS AND PACIFIERS

Hospitals support mothers to breastfeed by...



10 DISCHARGE

Hospitals support mothers to breastfeed by...





BREASTFEEDING | THE GOAL

By 2025, increase to at least 50% the rate of exclusive breastfeeding in the first six months

WHY IT MATTERS



Babies who are fed **nothing but breastmilk** from birth through their first 6 months of life get the **best start**



Exclusive breastfeeding provides babies: **the perfect nutrition** & everything they need for healthy growth and brain development



Protection from respiratory infections, diarrhoeal disease, and other **life-threatening ailments**



Protection against **obesity & non-communicable diseases** such as asthma and diabetes

Updated October 2018

RECOMMENDED ACTIONS

LIMIT FORMULA MARKETING

WHAT? Significantly limit the marketing of breastmilk substitutes



HOW? Strengthen the monitoring, enforcement and legislation related to the International Code of Marketing of Breastmilk Substitutes

SUPPORT PAID LEAVE

WHAT? Empower women to exclusively breastfeed



HOW? Enact six-months mandatory paid maternity leave and policies that encourage women to breastfeed in the workplace and in public

STRENGTHEN HEALTH SYSTEMS

WHAT? Provide hospital and health facilities-based capacity to support exclusive breastfeeding



HOW? Expand and institutionalize the baby-friendly hospital initiative in health systems

SUPPORT MOTHERS

WHAT? Provide community-based strategies to support exclusive breastfeeding counselling for pregnant and lactating women



HOW? Peer-to-peer and group counselling to improve exclusive breastfeeding rates, including the implementation of communication campaigns tailored to the local context

SCOPE OF THE PROBLEM


Globally, only **41% of infants** are exclusively breastfed



Suboptimal breastfeeding contributes to more than **800,000** infant deaths



Countries lose more than **\$300 billion** annually because of low breastfeeding rates



MCH in KSA

Maternal mortality in 1990-2015

WHO, UNICEF, UNFPA, World Bank Group, and United Nations Population Division
Maternal Mortality Estimation Inter-Agency Group

SAUDI ARABIA

Year	Maternal mortality ratio (MMR) ^a	Maternal deaths	AIDS-related indirect maternal deaths	Live births ^b	Proportion of maternal deaths among deaths of female reproductive age (PM %)
	Per 100 000 live births (lb)	Numbers	Numbers	Thousands	
1990	46 [32-67]	270	0	579	5.6
1995	33 [23-46]	190	0	581	4.2
2000	23 [16-34]	130	0	566	2.9
2005	18 [12-27]	100	0	578	2.3
2010	14 [8-23]	84	0	613	1.9
2015	12 [7-20]	72	0	619	1.6

^a MMR and PM are calculated for women 15-49 years.

^b Live birth data are from World Population Prospects: the 2015 Revision. New York, Population Division, Department of Economic and Social Affairs, United Nations Secretariat; 2015.

Annual Rate of Reduction (%)	
1990-2015	5.5 [3.7 - 7.5]
1990-2000	6.8 [4.2 - 9.6]
2000-2015	4.7 [2.3 - 7.1]
2005-2015	4.2 [1.4 - 7.1]

MCH Indicators in KSA

Under-5 mortality rank	141
Under-5 mortality rate (2012)	9
Infant Mortality rate per 1000 live births (under 1), (2012)	16.2
Annual rate of reduction (%) Under-5 mortality rate, (1990-2012)	7.7
Maternal mortality ratio (2010, adjusted)	24
Antenatal care coverage (%) at least 1 visit, 2008	97

- http://www.unicef.org/infobycountry/saudi-arabia_statistics.html, 2013
- Ministry of health KSA, 2012

ORIGINAL ARTICLE

Why mothers are not exclusively breast feeding their babies till 6 months of age? Knowledge and practices data from two large cities of the Kingdom of Saudi Arabia

Hafsa Raheel (1), Shabana Tharkar (2)

- (1) Department of Family and Community Medicine, College Of Medicine, King Saud University, Riyadh, Saudi Arabia
- (2) Prince Sattam Chair for Epidemiology and Public Health Research, Department of Family and Community Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia

ABSTRACT

The noble practice of breast feeding is on the decline across the globe. Our objective was to research why women stop feeding their infants before the recommended 6 months of exclusive breast feeding and to assess the mothers' knowledge regarding importance and benefits of breast feeding. A cross-sectional study was conducted in two cities of Riyadh and Dammam using a structured questionnaire to 614 Saudi females in reproductive age group (15–45 years) from February to April 2016. Majority of the respondents were <30 years old, housewives and multiparous. The breast feeding initiation rate was 76% while only 37% continued to exclusively breast feed the infants until 6 months. Mothers of Dammam city showed higher rates of overall breast feeding, higher initiation of breast feeding within 24 hours of delivery and longer duration of breast feeding practices. Mothers older than 30 years ($p < 0.014$), residents of Dammam

city ($P < 0.000$) and receiving information on breast feeding during antenatal care ($P < 0.001$) were associated with higher knowledge scores. Residents of Riyadh, working mothers, delayed initiation of breast feeding after 24 hours of giving birth, and those who did not get information on breast feeding during antenatal classes were at higher risk of stopping exclusive breast feeding before 6 months. Regional differences exist with Dammam city having greater awareness and better compliance to breast feeding practices. There is a need to strengthen the education facilities at ANC clinics in Riyadh regarding duration and benefits of breast feeding along with nationwide promotion of breast feeding practices as per guidelines.

KEYWORDS:

Breast feeding pattern; Knowledge; Barriers of breast feeding; Saudi Arabia.




The Necessary Vaccines for a Woman Before, During and After Pregnancy




Be sure to take the necessary vaccines before pregnancy in a timely manner to help maintain the health of the mother and the fetus, after consulting with the doctor

The Recommended Safe Vaccines



After Taking the Vaccine



Ready for pregnancy?

Iron

Add iron-rich foods to your diet to prevent anemia



Vaccinations

Make sure to get the necessary vaccines before getting pregnant



Folic acid Supplements

Take folic acid Supplements when Planning to get pregnant until week 12 of pregnancy



Vitamin D Supplements

Take vitamin D supplements for healthy teeth and bones

Consult your doctor before taking any medications or supplements



Nutrition during Pregnancy

Maintaining good nutrition and a healthy diet during pregnancy is critical for the health of the mother and fetus. It is also good to remember that the quality of the food is more important than the quantity, so it is recommended to choose healthy foods with high nutritional value.

Foods to avoid



Uncooked meat and eggs

May contain the Listeria bacteria that can be transmitted to the fetus through the placenta causing miscarriage, or stillbirth.



Raw fish and shellfish

May contain germs.



High mercury fish

Can damage the nervous system of the fetus (like: swordfish, and marlin)



Some types of oily fish

May contain harmful chemicals that accumulate in the body over time, (such as: Mackerel, sardines, salmon and fresh tuna).



Caffeine

Too much caffeine increases the risk of miscarriage and low birth weight.



Unpasteurized milk and all its products


May carry harmful bacteria that can cause diseases in the mother and fetus



How to Sleep During Pregnancy



Sleep on one of the 2 sides while supporting the belly with pillows



Sleeping on the back leads to a lack of blood and oxygen flow to the fetus



Make sure to relax before sleeping



Reduce your intake of tea, coffee and soft drinks before sleep



Make sure to nap



Place a pillow between your knees



وزارة الصحة
Ministry of Health

Pregnant Woman and Exercise



It prepare the body for giving birth



You can exercise after consulting your doctor



Contributes to prevention of back pain and constipation



Helps adapt to physical and mental changes



Best Kinds of Exercise



Swimming



Walking



Yoga




وزارة الصحة
Ministry of Health

The Myth of the Narrow Pelvis

✦ **The female pelvis is not made up of one solid bone;**

it consists of many bones and ligaments that loosen and move during childbirth




Having a small pelvis

does not prevent women from giving birth naturally




The Signs of True Labor



What is the Normal Weight Gain during Pregnancy?

If the fetus weighs between 3 kg to 3.5 kg



the total weight gain in terms of the natural increase of weight during pregnancy



Caesarean sections should only be performed when medically necessary



If your doctor determines that a natural birth is no longer possible even with some medical intervention, then you will need an emergency caesarean section.

Postpartum Depression

Has no specific cause, it may result from several physical or psychological factors involving changes in



Lack of sleep and stress



Changes in thyroid hormones



Changes in female hormones

Risk Factors for Postpartum Depression



Personal history of mental illness



Family history of mental illness



Lack of moral support from family and friends



Having faced problems with previous pregnancy



Anxiety and stress

Why are Postpartum Checkups Important?



To discuss required vaccines



To monitor the mother's blood pressure especially if she faced any issues during or after delivery



To examine the incision position and the stitches in case of caesarean section or any surgical intervention



To discuss how to regulate the menstrual cycle



To discuss vaginal discharge, especially if it is persistent



To discuss possible contraception methods



To monitor the mother's health and recovery



To discuss the mother's general emotional state

It is recommended to schedule a postpartum checkup 6 weeks after delivery

References

- *Lale Say et al. Global causes of maternal death: a WHO systematic analysis. The Lancet Global Health .Volume 2, Issue 6, Pages e323-e333 (June 2014) . DOI: 10.1016/S2214-109X(14)70227-X*
- Levels & Trends in Child Mortality Report 2015 Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation. United Nations Available at: <http://www.childmortality.org/>
- Mohammad Afzal Mahmood, Hafsa Raheel, et al., “Root-Cause Analysis of Persistently High Maternal Mortality in a Rural District of Indonesia: Role of Clinical Care Quality and Health Services Organizational Factors,” *BioMed Research International*, vol. 2018, Article ID 3673265, 11 pages, 2018. doi:10.1155/2018/3673265
- **Success factors for reducing maternal and child mortality. Policy and practice, WHO. Available at: <http://www.who.int/bulletin/volumes/92/7/14-138131/en/>**