



Tutorial 7: Vaccination/Immuni- zation (counseling)



Color index:

- Main text
- Males slides
- Females slides
- Doctor notes
- Important
- Extra

Vaccination/Immunization (counseling)

01

A mother came to your clinic with her 4 month old baby for the well-baby clinic, he is due for vaccination. His pre, peri and postnatal history are not significant. On examination: He is febrile T.38 °C, mild throat congestion and clear chest. Other examinations are normal. How will you proceed with the consultation?

We will give him the vaccine because in mild temperature it is not contraindicated

02

9 months-old-girl was brought by her parents to your clinic with cough, wheezy breathing and runny nose for 2 days. O/E T.38.1°C, congested throat, chest, Occasional wheeze. The Child is active and feeding well. Parents are worried about giving vaccination to their child, how are you going to deal with the parents, and will you vaccinate the child? The difference from case 1 is that the baby has URT symptoms, is it now contraindicated to vaccinate?

NO, not contraindicated. Give her the vaccine

03

A parents are concerned about MMR (Measles, Mumps, and Rubella) vaccine and that it cause autism, as they read something about that. What would you tell them?

We educate and ensure the parents that there's no relationship between autism and the vaccine.

04

A child was brought by his parents at the age of 9 months. The last vaccine he received was at the age of 4 months. What will you do? (That means there is a skipped vaccine at the age of 6 months) Do we repeat the schedule or just give him the ones he missed?

We give both vaccines (the one at 6 & 9) and reassure as well as counsel the parents to make sure it doesn't happen again. We continue as it is, no need to restart because of the delay.

لو نفس الحالة وراح المريض ولقى بالمستشفى بس فاكسين واحد وقالوا له المستشفى الفاكسين الثاني بيجي بكرا وش راح تسوي؟ تعطيه الفاكسين اللي موجود اليوم ، والثاني بعد ٤ اسابيع (يصير بينهم فترة ٤ اسابيع) ليه؟ عشان ما يتلخبط الجسم ويقال تكوين Antibodies

Bottom line: Either you give the two live vaccines together or 4 weeks apart

Vaccination/Immunization (counseling)

05

A concerned mother consult you about her 1 year old child, she is not sure if he had received BCG vaccine. Which arm you are going to examine in the child and why?
Left arm. BCG causes necrosis so it leaves a scar like trace and having a uniform location makes it easier to find it. According to Dr. Abdullah if you don't find it you should give the vaccine. According to Dr. Noura if not found we usually don't give it , we check the vaccination booklet of the child. Even if you do not find it there you do not give the vaccine. In KSA the birth certificate isn't given unless taken but now it has been postponed to 6 months

BCG vaccine : Left upper arm

Measles vaccine: Right upper arm

06

A child is due for MMR vaccine at the age of 12 months. The mother asks you about the side effects. What are you going to tell her?

Localized pain at the injection site fever 3-5 days

Mild form of the disease (parotid gland inflammation)

07

A child was given the first two doses of hepatitis vaccine but he skipped the third (he didn't take it on date) will you repeat from the start of continue? We will continue.

Click here to go to the [immunization schedule](#)

Team leaders

Alaa Alsulmi

Abdulaziz Alghuligah

Khaled Alsubaie

Done by
Gaida Alassery

