


# COPD

Dr R Nadama MD MRCP(lond) MRCP(UK), FRCP(Lond), EDARM, FCCP

# Definition

Chronic Obstructive Pulmonary Disease (COPD) is a common, preventable and treatable disease that is characterized by persistent respiratory symptoms and airflow limitation that is due to airway and/or alveolar abnormalities usually caused by significant exposure to noxious particles or gases.

**Figure 1.1. Etiology, pathobiology and pathology of COPD leading to airflow limitation and clinical manifestations**



# Chronic Obstructive Pulmonary Disease (COPD)

Characterized by respiratory symptoms and airflow limitation that is due to airway and/or alveolar abnormalities usually caused by significant exposure to noxious particles or gases.

The main risk factor for COPD is tobacco smoking but other environmental exposures such as biomass fuel exposure and air pollution may contribute

# Prevalence

Prevalence of COPD was higher in smokers and ex-smokers compared to non-smokers

Higher  $\geq 40$  year group compared to those  $< 40$

Higher in men than women

# Prevalence

- Estimated 384 million COPD cases in 2010.
- Three million deaths annually.
- Increasing smoking
- By 2030 predicted 4.5 million COPD related deaths annually

# Pathology, pathogenesis & pathophysiology

## Pathology

- Chronic inflammation
- Structural changes


## Pathogenesis

- Oxidative stress
- Protease-antiprotease imbalance
- Inflammatory cells
- Inflammatory mediators
- Peribronchiolar and interstitial fibrosis

## Pathophysiology

- Airflow limitation and gas trapping
- Gas exchange abnormalities
- Mucus hypersecretion
- Pulmonary hypertension

# Pathogenesis




# Diagnosis and Initial Assessment

# Diagnosis


Figure 2.1. Pathways to the diagnosis of COPD


---



# Symptoms

- Chronic and progressive dyspnea
- Cough
- Sputum production
- Wheezing and chest tightness
- Others – including fatigue, weight loss, anorexia, syncope, rib fractures, ankle swelling, depression, anxiety.





**Table 2.2. Other causes of chronic cough**

**Intrathoracic**

- Asthma
- Lung cancer
- Tuberculosis
- Bronchiectasis
- Left heart failure
- Interstitial lung disease
- Cystic fibrosis
- Idiopathic cough

**Extrathoracic**

- Chronic allergic rhinitis
- Post nasal drip syndrome (PNDS)
- Upper Airway Cough Syndrome (UACS)
- Gastroesophageal reflux
- Medication (e.g. ACE inhibitors)

# Spirometry

**Table 2.4. Classification of airflow limitation severity in COPD (Based on post-bronchodilator FEV<sub>1</sub>)**

In patients with FEV<sub>1</sub>/FVC < 0.70:

GOLD 1:	Mild	FEV <sub>1</sub> ≥ 80% predicted
GOLD 2:	Moderate	50% ≤ FEV <sub>1</sub> < 80% predicted
GOLD 3:	Severe	30% ≤ FEV <sub>1</sub> < 50% predicted
GOLD 4:	Very Severe	FEV <sub>1</sub> < 30% predicted

# Choice of thresholds

- COPD Assessment Test (CAT TM )
- Modified Medical Research Council (mMRC) questionnaire

Figure 2.3. CAT Assessment

For each item below, place a mark (X) in the box that best describes you currently. Be sure to only select one response for each question.

**Example:** I am very happy ①  ② ③ ④ ⑤ I am very sad

	SCORE
I never cough ① ② ③ ④ ⑤ I cough all the time	<input type="checkbox"/>
I have no phlegm (mucus) in my chest at all ① ② ③ ④ ⑤ My chest is completely full of phlegm (mucus)	<input type="checkbox"/>
My chest does not feel tight at all ① ② ③ ④ ⑤ My chest feels very tight	<input type="checkbox"/>
When I walk up a hill or one flight of stairs I am not breathless ① ② ③ ④ ⑤ When I walk up a hill or one flight of stairs I am very breathless	<input type="checkbox"/>
I am not limited doing any activities at home ① ② ③ ④ ⑤ I am very limited doing activities at home	<input type="checkbox"/>
I am confident leaving my home despite my lung condition ① ② ③ ④ ⑤ I am not at all confident leaving my home because of my lung condition	<input type="checkbox"/>
I sleep soundly ① ② ③ ④ ⑤ I don't sleep soundly because of my lung condition	<input type="checkbox"/>
I have lots of energy ① ② ③ ④ ⑤ I have no energy at all	<input type="checkbox"/>
<b>TOTAL SCORE</b>	<input type="checkbox"/>


Reference: Jones et al. ERJ 2009; 34 (3); 648-54.

Table 2.5. Modified MRC dyspnea scale <sup>a</sup>	
PLEASE TICK IN THE BOX THAT APPLIES TO YOU (ONE BOX ONLY) (Grades 0-4)	
mMRC Grade 0. I only get breathless with strenuous exercise.	<input type="checkbox"/>
mMRC Grade 1. I get short of breath when hurrying on the level or walking up a slight hill.	<input type="checkbox"/>
mMRC Grade 2. I walk slower than people of the same age on the level because of breathlessness, or I have to stop for breath when walking on my own pace on the level.	<input type="checkbox"/>
mMRC Grade 3. I stop for breath after walking about 100 meters or after a few minutes on the level.	<input type="checkbox"/>
mMRC Grade 4. I am too breathless to leave the house or I am breathless when dressing or undressing.	<input type="checkbox"/>

<sup>a</sup> Fletcher CM. BMJ 1960; 2: 1662.

# ABCD of COPD

Figure 2.4. The refined ABCD assessment tool



# Alpha-1 antitrypsin deficiency (AATD)

- The World Health Organization recommends that all patients with a diagnosis of COPD should be screened once especially in areas with high AATD prevalence.
- AATD patients are typically < 45 years with panlobular basal emphysema
- Delay in diagnosis in older AATD patients presents as more typical distribution of emphysema (centrilobular apical).
- A low concentration (< 20% normal) is highly suggestive of homozygous deficiency.

# Differential Diagnosis

# Management

Approach

# Management of Stable COPD

Once COPD has been diagnosed, effective management should be based on an individualized assessment to reduce both current symptoms and future risks of exacerbations.

**Table 4.1. Goals for treatment of stable COPD**

- Relieve symptoms
  - Improve exercise tolerance
  - Improve health status
- 
- Diagram: A vertical bracket on the right side of the three bullet points above groups them, with an arrow pointing to the text 'REDUCE SYMPTOMS'.

*and*

- Prevent disease progression
  - Prevent and treat exacerbations
  - Reduce mortality
- 
- Diagram: A vertical bracket on the right side of the three bullet points above groups them, with an arrow pointing to the text 'REDUCE RISK'.

# Non-Pharmacologic Treatment

- Education and self-management
- Physical activity
- Pulmonary rehabilitation programs
- Exercise training
- Self-management education
- End of life and palliative care
- Nutritional support
- Vaccination
- Oxygen therapy

# Smoking

- Smoking cessation has the greatest capacity to influence the natural history of COPD.
- If effective resources and time are dedicated to smoking cessation, long-term quit success rates of up to 25% can be achieved.




FIGURE 2 - Risks for various men if they smoke: differences between these lines illustrate effects that smoking, and stopping smoking can have on FEV<sub>1</sub> of man who is liable to develop Chronic obstructive lung disease if he smokes. (BMJ, 1977)

# Vaccination

- Influenza vaccination can reduce serious illness (such as lower respiratory tract infections requiring hospitalization)<sup>24</sup> and death in COPD patients.
- Pneumococcal vaccinations, PCV13 and PPSV23, are recommended for all patients  $\geq 65$  years of age

**Table 3.2. Vaccination for stable COPD**

- Influenza vaccination reduces serious illness and death in COPD patients (**Evidence B**).
- The 23-valent pneumococcal polysaccharide vaccine (PPSV23) has been shown to reduce the incidence of community-acquired pneumonia in COPD patients aged  $< 65$  years with an  $FEV_1 < 40\%$  predicted and in those with comorbidities (**Evidence B**).
- In the general population of adults  $\geq 65$  years the 13-valent conjugated pneumococcal vaccine (PCV13) has demonstrated significant efficacy in reducing bacteremia and serious invasive pneumococcal disease (**Evidence B**).

# Pulmonary Rehabilitation

## **Exercise training**

– 6–12 weeks;

longer programs result in larger effects

– 20–30 min walking per session, to limits of symptoms

## **Patient education**

- Smoking cessation,
- COPD natural history and management,
- self-management,
- exacerbations

## **Assessment and follow-up**

## **Nutritional support**

# Rehabilitation, Education & Self-Management

**Table 3.8. Pulmonary rehabilitation, self-management and integrative care in COPD**

## **Pulmonary rehabilitation**

- Pulmonary rehabilitation improves dyspnea, health status and exercise tolerance in stable patients (**Evidence A**).
- Pulmonary rehabilitation reduces hospitalizations among patients who have had a recent exacerbation ( $\leq 4$  weeks from prior hospitalization) (**Evidence B**).

## **Education and self-management**

- Education alone has not been shown to be effective (**Evidence C**).
- Self-management intervention with communication with a health care professional improves health status and decreases hospitalizations and emergency department visits (**Evidence B**).

## **Integrated care programs**

- Integrated care and telehealth have no demonstrated benefit at this time (**Evidence B**).

# Pharmacological therapy

Table 3.3. Commonly used maintenance medications in COPD*					
Drug	Inhaler (mcg)	Solution for nebulizer (mg/ml)	Oral	Vials for injection (mg)	Duration of action (hours)
<b>Beta<sub>2</sub>-agonists</b>					
<i>Short-acting</i>					
Fenoterol	100-200 (MDI)	1	2.5 mg (pill), 0.05% (syrup)		4-6
Levalbuterol	45-90 (MDI)	0.1, 0.21, 0.25, 0.42			6-8
Salbutamol (albuterol)	90, 100, 200 (MDI & DPI) <sup>†</sup>	1, 2, 2.5, 5 mg/ml	2, 4, 5 mg (pill), 8 mg (extended release tablet) 0.024%/0.4 mg (syrup)	0.1, 0.5 mg	4-6, 12 (ex- tended release)
Terbutaline	500 (DPI)		2.5, 5 mg (pill)	0.2, 0.25, 1 mg	4-6
<i>Long-acting</i>					
Arformoterol		0.0075 <sup>†</sup>			12
Formoterol	4.5-9 (DPI)	0.01 <sup>^</sup>			12
Indacaterol	75-300 (DPI)				24
Olodaterol	2.5, 5 (SMI)				24
Salmeterol	25-50 (MDI & DPI)				12
<b>Anticholinergics</b>					
<i>Short-acting</i>					
Ipratropium bromide	20, 40 (MDI)	0.2			6-8
Oxipropium bromide	100 (MDI)				7-9
<i>Long-acting</i>					
Acidinium bromide	400 (DPI), 400 (MDI)				12
Glycopyrronium bromide	15.6 & 50 (DPI) <sup>†</sup>		1 mg (solution)	0.2 mg	12-24
Tiotropium	18 (DPI), 2.5 & 5 (SMI)				24
Umeclidinium	62.5 (DPI)				24
<b>Combination of short-acting beta<sub>2</sub>-agonist plus anticholinergic in one device</b>					
Fenoterol/ipratropium	50/20 (SMI)	1.25, 0.5 mg in 4ml			6-8
Salbutamol/ipratropium	100/20 (SMI), 75/15 (MDI)	0.5, 2.5 mg in 3ml			6-8

# Pharmacologic Therapy

Combination of long-acting beta <sub>2</sub> -agonist plus anticholinergic in one device			
Formoterol/aclidinium	12/400 (DPI)		12
Formoterol/glycopyrronium	9.6/18 (MDI)		12
Indacaterol/glycopyrronium	27.5/15.6 & 110/50 (DPI) <sup>†</sup>		12-24
Vilanterol/umeclidinium	25/62.5 (DPI)		24
Olodaterol/tiotropium	5/5 (SMI)		24
Methylxanthines			
Aminophylline	105 mg/ml (solution)	250, 500 mg	Variable, up to 24
Theophylline (SR)	100-600 mg (pill)	250, 400, 500 mg	Variable, up to 24
Combination of long-acting beta <sub>2</sub> -agonist plus corticosteroids in one device			
Formoterol/beclomethasone	6/100 (MDI)		
Formoterol/budesonide	4.5/160 (MDI), 4.5/80 (MDI), 9/320 (DPI), 9/160 (DPI)		
Formoterol/mometasone	10/200, 10/400 (MDI)		
Salmeterol/fluticasone	5/100, 50/250, 5/500 (DPI), 21/45, 21/115, 21/230 (MDI)		
Vilanterol/fluticasone furoate	25/100 (DPI)		
Phosphodiesterase-4 inhibitors			
Roflumilast		500 mcg (pill)	

MDI = metered dose inhaler; DPI = dry powder inhaler; SMI = soft mist inhaler

\* Not all formulations are available in all countries; in some countries other formulations and dosages may be available

<sup>†</sup> Dose availability varies by country

<sup>‡</sup> Formoterol nebulized solution is based on the unit dose vial containing 20 mcg in a volume of 2.0 ml

<sup>§</sup> Dose varies by country

# Anti-inflammatory Therapy in Stable COPD

**Table 3.5. Anti-inflammatory therapy in stable COPD**

## Inhaled corticosteroids

- An ICS combined with a LABA is more effective than the individual components in improving lung function and health status and reducing exacerbations in patients with exacerbations and moderate to very severe COPD (**Evidence A**).
- Regular treatment with ICS increases the risk of pneumonia especially in those with severe disease (**Evidence A**).
- Triple inhaled therapy of ICS/LAMA/LABA improves lung function, symptoms and health status (**Evidence A**) and reduces exacerbations (**Evidence B**) compared to ICS/LABA or LAMA monotherapy.

## Oral glucocorticoids

- Long-term use of oral glucocorticoids has numerous side effects (**Evidence A**) with no evidence of benefits (**Evidence C**).

## PDE4 inhibitors

- In patients with chronic bronchitis, severe to very severe COPD and a history of exacerbations:
  - » A PDE4 inhibitor improves lung function and reduces moderate and severe exacerbations (**Evidence A**).
  - » A PDE4 inhibitor improves lung function and decreases exacerbations in patients who are on fixed-dose LABA/ICS combinations (**Evidence B**).

## Antibiotics

- Long-term azithromycin and erythromycin therapy reduces exacerbations over one year (**Evidence A**).
- Treatment with azithromycin is associated with an increased incidence of bacterial resistance (**Evidence A**) and hearing test impairments (**Evidence B**).

## Mucolytics/antioxidants

- Regular use of NAC and carbocysteine reduces the risk of exacerbations in select populations (**Evidence B**).

## Other anti-inflammatory agents


- Simvastatin does not prevent exacerbations in COPD patients at increased risk of exacerbations and without indications for statin therapy (**Evidence A**). However, observational studies suggest that statins may have positive effects on some outcomes in patients with COPD who receive them for cardiovascular and metabolic indications (**Evidence C**).
- Leukotriene modifiers have not been tested adequately in COPD patients.

# The Inhaled Route

Right inhaler for the Right patient

# ABCD of COPD

Figure 2.4. The refined ABCD assessment tool





# Oxygen therapy

Long-term oxygen therapy is indicated for stable patients who have:

- PaO<sub>2</sub> at or below 7.3 kPa (55 mmHg) or SaO<sub>2</sub> at or below 88%, with or without hypercapnia confirmed twice over a three week period; or
- PaO<sub>2</sub> between 7.3 kPa (55 mmHg) and 8.0 kPa (60 mmHg), or SaO<sub>2</sub> of 88%, if there is evidence of pulmonary hypertension, peripheral edema suggesting congestive cardiac failure, or polycythemia (hematocrit > 55%).

# Interventional bronchoscopy and surgery

Surgical or endoscopic LVRS

Bullectomy

Transplant

# Management of Exacerbations

**COPD exacerbations** are defined as an acute worsening of respiratory symptoms that result in additional therapy.

- They are classified as:
  - Mild (treated with short acting bronchodilators only, SABDs)
  - Moderate (treated with SABDs plus antibiotics and/or oral corticosteroids) or
  - Severe (patient requires hospitalization or visits the emergency room). Severe exacerbations may also be associated with acute respiratory failure.

# Classification of hospitalized patients

## ***No respiratory failure:***

- Respiratory rate: 20-30 breaths per minute; no use of accessory respiratory muscles; no changes in mental status; hypoxemia improved with supplemental oxygen given via Venturi mask 28-35% inspired oxygen ( $\text{FiO}_2$ ); no increase in  $\text{PaCO}_2$ .

## ***Acute respiratory failure — non-life-threatening:***

- Respiratory rate: > 30 breaths per minute; using accessory respiratory muscles; no change in mental status; hypoxemia improved with supplemental oxygen via Venturi mask 25-30%  $\text{FiO}_2$ ; hypercarbia i.e.,  $\text{PaCO}_2$  increased compared with baseline or elevated 50-60 mmHg

## ***Acute respiratory failure — life-threatening:***

- Respiratory rate: > 30 breaths per minute; using accessory respiratory muscles; acute changes in mental status; hypoxemia not improved with supplemental oxygen via Venturi mask or requiring  $\text{FiO}_2 > 40\%$ ; hypercarbia i.e.,  $\text{PaCO}_2$  increased compared with baseline or elevated > 60 mmHg or the presence of acidosis ( $\text{pH} \leq 7.25$ ).

# Management of Exacerbations

- Bronchodilators
- Corticosteroids
- Antibiotics
- Controlled oxygen therapy
- Ventilatory support; non invasive/invasive

# Management of Exacerbations

## NIV

### Table 5.5. Indications for noninvasive mechanical ventilation (NIV)

At least one of the following:

- Respiratory acidosis ( $\text{PaCO}_2 \geq 6.0$  kPa or 45 mmHg and arterial pH  $\leq 7.35$ ).
- Severe dyspnea with clinical signs suggestive of respiratory muscle fatigue, increased work of breathing, or both, such as use of respiratory accessory muscles, paradoxical motion of the abdomen, or retraction of the intercostal spaces.
- Persistent hypoxemia despite supplemental oxygen therapy.

# Management of Exacerbations

## Respiratory support

### Table 5.4. Indications for respiratory or medical intensive care unit admission\*

- Severe dyspnea that responds inadequately to initial emergency therapy.
- Changes in mental status (confusion, lethargy, coma).
- Persistent or worsening hypoxemia ( $\text{PaO}_2 < 5.3 \text{ kPa}$  or  $40 \text{ mmHg}$ ) and/or severe/worsening respiratory acidosis ( $\text{pH} < 7.25$ ) despite supplemental oxygen and noninvasive ventilation.
- Need for invasive mechanical ventilation.
- Hemodynamic instability—need for vasopressors.

\*Local resources need to be considered.

