




Editing file

Arrhythmias



Objectives :

- ★ General approach to arrhythmias
- ★ Specific types and chronic management:
 - Atrial Fibrillation (AF)
 - Atrial Flutter (AFL)
 - Supraventricular tachycardia (SVT)
 - Wolf-Parkinson-White syndrome (WPW)
 - Ventricular tachycardia (VT)
 - Ventricular fibrillation (VF)
- ★ Acute management

Color index

- Original text
- Females slides
- Males slides
- Doctor's notes ⁴³⁸
- Doctor's notes ⁴³⁹
- Text book
- Important
- Golden notes
- Extra




Normal cardiac rhythm (Sinus rhythm)

- **Characteristics of normal cardiac rhythm:**
 - **Heart rate (60-100 bpm)**
 - Abnormal: Check table
 - **Origin of impulses should be from S.A node (sinus node)**
 - Abnormal: Any place other than sinus node e.g.
 - **Supraventricular arrhythmias:**¹ AV node (Nodal/junctional arrhythmias), Atrium it self (Atrial arrhythmias)
 - **Ventricular arrhythmias**
 - **Cardiac impulse should propagate through normal conduction pathway***
 - **Normal velocity**

> 350 bpm	Fibrillation
250-350 bpm	Flutter
150-250 bpm	Paroxysmal tachycardia
100-150 bpm	Simple Tachycardia
60-100 bpm	Sinus rhythm
< 60 bpm	Bradycardia

* What's the normal conduction pathway?

Depolarisation **starts in the sinoatrial node** and spreads through the atria (**blue arrows**), and then through the **atrioventricular node (black arrows)**. Depolarisation then spreads through the **bundle of His** and the **bundle branches** and **purkinje fibres** to reach the ventricular muscle (**red arrows**). Repolarisation spreads from epicardium to endocardium (**green arrows**).



Action potential of cardiac muscles


Phase 0: Depolarization occurs through fast Na⁺ channels.

Phase 1: Na⁺ channels close and K⁺ starts leaving the cell, causing slight repolarization.

Phase 2: or better known as the “plateau phase”, Ca²⁺ gets in the cell and K⁺ leaves the cell


Phase 3: K⁺ leaves the cell → restores resting membrane potential RMP

Phase 4: maintained by Na⁺/K⁺ ATPase channel



ECG basics

- **P wave:** Atrial depolarisation
- **QRS complex:** ventricular myocardial depolarisation.
- **T wave:** For ventricular repolarization, while Atrial repolarization does not cause a detectable signal
- **PR interval:** it largely reflects the duration of AV nodal conduction.
- **QT interval:** represents the total duration of ventricular depolarisation and repolarisation.
- **RR interval:** Time between beats - is used to calculate heart rate



¹- Sinus arrhythmias are also considered supraventricular arrhythmias, but it differs from the other two in that it's origin is from S.A node.

General approach to arrhythmias

History :

HPI	Palpitation ¹ , Dizziness , Syncope ² , fatigue, chest pain and SOB ³ , stroke (AF and atrial flutter)
Family history	
Social history	<ul style="list-style-type: none"> • Ethol, illicit drug use
Past medical history (PMH)	<ul style="list-style-type: none"> • Underlying heart disease • Past medical and surgical history

Work-up :

- ECG (baseline ⁴ and **during episodes**), Holter ⁵ , Echo ⁶ (explained in next slides)
- R/out secondary causes: electrolytes, TSH, sleep study

Management : in general :

- Wait and see ⁷
- Medical therapy
- Ablation ⁸ كي القلب



- Palpitation is when you feel your heart beats (doesn't have to be fast or irregular). It could be :
 - normal : while doing exercise (extensive workout)
 - abnormal : eg, at rest
- Definition: sudden , transient loss of consciousness .
- Especially in patient with underlying heart disease .
- Gives clues about the type of the arrhythmia .
- If the arrhythmia happen infrequently .
- To make sure that there is no underlying heart disease .
- When you get one episode of arrhythmia every couple of years .
- The insertion of a catheter through the groin all the way up to the heart then use radiofrequency to burn the focus



General info



- **Atrial fibrillation (AF)** is the **most common** sustained cardiac arrhythmia, with prevalence **rising with age (>55y/o)**. It's also **more common in males**.
- It is associated with significant morbidity and a two-fold increase in mortality.
- **AF is associated with a five-fold increased risk of stroke**, primarily as a result of embolism of a thrombus that has formed in the atrium

AF is the most common overall (30-40%) followed by Ventricular tachycardia (VT) then AV block then Ventricular fibrillation (VF) then Sick sinus syndrome (SSS) then Supraventricular tachycardia (SVT)
Other less common arrhythmias: ventricular premature contraction (VPC), Premature atrial contraction (PAC), Long QTs syndrome (LQTS)¹ and Brugada syndrome²



It shows the current and the estimated prevalence of AF; as you can see the prevalence of AF is increasing in the future because the risk factors (**the most important is the age**) are common and they will be more common in the future


Men have higher incidence in all age strata
Saudi Arabia is in the green zone due to the young population



AF is more common as you age

Pathogenesis

- AF is a complex arrhythmia characterised by both abnormal automatic firing and the presence of **multiple interacting re-entry microcircuits looping around the atria**, most often located within pulmonary veins.
- During episodes of AF, the atria beat rapidly but in an uncoordinated, **chaotic** and **ineffective manner** → ↓ Cardiac Output
- The AV node conducts a proportion of the atrial impulses to produce an irregular ventricular response, giving rise to an **irregularly irregular pulse**.
- Loss of atrial contraction and left atrial dilatation cause stasis of blood in the LA and may lead to **thrombus formation in the left atrial appendage**. This predisposes patients to **stroke** and other forms of systemic embolism.



Risk factors

The most important ones are the highlighted

- **Obesity**
- **Hypertension**
- **Alcohol consumption**
- **Obstructive sleep apnea**
- Smoking
- Lipid profile
- (Pre-) Diabetes
- Vascular disease
- Coronary artery disease
- HF
- Physical inactivity
- Chronic kidney disease
- COPD
- Valvular disease
- Inflammatory disease



1- A congenital disorder characterized by a prolongation of the QT interval on ECG and a propensity to ventricular tachyarrhythmias, which may lead to syncope, cardiac arrest or sudden death.
2- A genetically inherited disorder that causes a repolarization abnormality in cardiac myocytes that can lead to sudden cardiac death, especially in athletes and young people. It is characterized by ECG findings including the presence of a "pseudo-right bundle branch block" and persisting ST elevations in V1-V2.

Causes of AF **3 PARTIES**

Cardiac causes	Non-cardiac causes
<ul style="list-style-type: none"> • Most common causes: Hypertensive heart disease & Heart failure • Ischemic heart disease • Valvular heart disease: <ul style="list-style-type: none"> ○ Rheumatic (major cause in KSA): mitral stenosis ○ Non-rheumatic: aortic stenosis, mitral regurgitation • Pericarditis • Cardiac tumors: Atrial myxoma • Sick sinus syndrome¹ (Tachy-Brady syndrome) • Cardiomyopathy: <ul style="list-style-type: none"> ○ Hypertrophic ○ Idiopathic dilated (? cause vs. effect) • Post-coronary bypass surgery: 50% post cardiac patients develop AF 	<ul style="list-style-type: none"> • Pulmonary: <ul style="list-style-type: none"> ○ COPD ○ Pneumonia ○ Pulmonary embolism ★ Metabolic: <ul style="list-style-type: none"> ○ Throid disease: hyperthyroidism ○ Electrolyte disorder • Toxic: <ul style="list-style-type: none"> ○ Alcohol "Ethanol" ('holiday heart' syndrome)

Heart rhythm in AF

Normal heart rhythm is disrupted in AF

- **Rapid** (350–600 beats/min)² and **irregular atrial rhythm**.
- **Reduced filling** of the left and right ventricles.

Conduction of most impulses from the atria to ventricles is **blocked at the AV node**

Contraction of the ventricles can be:

- **Fast AF:** Irregular and rapid (110–180 beats/min; tachycardia), **Doesn't exceed 180bpm because of AV node delay.**
- **Slow AF:** Irregular and slow (<50 beats/min; bradycardia) **seen in elderly (>70y/o)**³
- **Controlled AF:** Normal

Cardiac output can be reduced
(If AV node fails to block the impulses from atria)

1- Due to wear and tear process in SA node. Sick sinus syndrome (SSS) refers to the dysfunction of the sinoatrial node and is responsible for several types of arrhythmia. It comprises bradyarrhythmias (e.g., sinus bradycardia, sinoatrial pauses, blocks, and arrest), and may alternate with supraventricular tachyarrhythmias, in which case it is referred to as tachycardia-bradycardia syndrome.



2- Will the AV node deliver all these beats? No, there's AV node delay. It has a feature called **decremental conduction**, which will automatically block its conduction once it senses it is over conducting to give the ventricles time to fill. Recall $CO = SV \times HR$, If HR is very high the ventricles will not have time to fill → Decrease in preload → Decrease SV → Decreased CO.

3- Because of AV node degeneration (wear and tear), the AV node **حيل ما فيها حيل** to conduct.

Atrial fibrillation (AF) ECG



ECG changes in AF

Normal ECG



- Normal heart rate (60-100 bpm)
- Regular rhythm
- Normal P Waves and QRS complexes
- Steady baseline


★ AF ECG



1. **Absent P waves** (Because atria doesn't contract), only a fine oscillation of the baseline (so-called fibrillation or f waves)
2. **Fast and irregular ventricular response (QRS complex)** → **irregular RR intervals (irregularly irregular rhythm)^{1&2}**
3. **Increased heart rate (Tachyarrhythmias)**
4. **Irregular baseline**

★ How to recognize AF in physical examination ?


If you feel the pulse and is irregularly irregular



1. There are 2 types of irregularity:
 - a. Regularly irregular: Cyclic irregularity which means it's predictable and has a pattern → Atrial flutter
 - b. Irregularly irregular: Not predictable and has no pattern → Atrial fibrillation
2. Also in physical examination if the pulse is irregularly irregular > most likely AF

◀ AF Begets (Generates) AF

- AF gives rise to AF, how?.




Over time AF tends to become persistent or permanent. However, Many episodes of AF resolve spontaneously.

◀ Signs & Symptoms

- The typical presentation is **with palpitation, breathlessness and fatigue**.
- It's usually **episodic** with symptoms being absent or intermittent
- **Up to 90% of episodes may not cause symptoms**
- Symptoms vary according to: Irregularity and rate of ventricular response, Functional status, AF duration, Patient factors, Comorbidities.
- **Patient may present with or without detectable heart diseases.**

Cause	Sign / Symptom
Irregular heart beat	<ul style="list-style-type: none"> • Irregularly irregular pulse • Palpitations
Cardiac ischemia	<ul style="list-style-type: none"> • Chest pain (Angina)
Hypotension	<ul style="list-style-type: none"> • Dizziness and fainting (syncope)
Increased risk of clot formation	<ul style="list-style-type: none"> • Thromboembolic TIA → Stroke Could be the first presentation!
Decreased Cardiac output	<ul style="list-style-type: none"> • Fatigue • Diminished exercise capacity • Breathlessness (dyspnoea) • Weakness (asthenia)



◀ Signs & Symptoms cont'

European Heart Rhythm Association (EHRA) Score:

Score	Symptoms	Description
1	None	AF does not cause any symptoms
2a	mild	Normal daily activity not affected by symptoms related to AF
2b	Moderate	Normal daily activity not affected by symptoms related to AF, but patient troubled by symptoms
3	Severe	Normal daily activity affected by symptoms related to AF
4	Disabling	Normal daily activity discontinued

◀ Classification of AF

Class	Definition
Paroxysmal AF	<ul style="list-style-type: none"> AF that terminates spontaneously or with intervention within 7 d of onset. Episodes may recur with variable frequency.
Persistent AF	<ul style="list-style-type: none"> Continuous AF that is sustained > 7 d but less than 12 months.
Long-standing persistent AF	<ul style="list-style-type: none"> Continuous AF >12 months in duration.
Permanent AF	<ul style="list-style-type: none"> The term “permanent AF” is used when the patient and clinician make a joint decision to stop further attempts to restore and/or maintain sinus rhythm. Acceptance of AF represents a therapeutic attitude on the part of the patient and clinician rather than an inherent pathophysiological attribute of AF. Acceptance of AF may change as symptoms, efficacy of therapeutic interventions, and patient and clinician preferences evolve
Lone or primary	<ul style="list-style-type: none"> Presence of paroxysmal, persistent or permanent atrial fibrillation with no evidence of cardiopulmonary or structural heart disease. Patients with lone AF are usually <60 and have less risk for thromboembolism.
Non Valvular AF	<ul style="list-style-type: none"> AF in the absence of rheumatic mitral stenosis, a mechanical or bioprosthetic heart valve, or mitral valve repair. No longer used

1. History and physical examination

01



Clinical conditions associated with AF

- **Underlying heart conditions** (e.g. valvular heart disease, heart failure, coronary artery disease, hypertension)
- Other **reversible conditions**
- **Ask about possible causes (Discussed earlier)**

02



Family history

- **Familial AF** (lone AF in a family)
- **AF secondary** to other **genetic conditions** (familial cardiomyopathies)

03



Type of AF

- **First episode, paroxysmal, persistent, permanent**
- **Triggers** – e.g. emotional stress, alcohol, physical exercise, gastroesophageal disease
- **Specific symptoms**
- **Response to any treatments** administered

2. ECG

- Essential for **ALL** patients with AF to identify:



1

Abnormal
heart
rhythm¹
(verify AF)

2

Pre-
excitation

3

Prior MI

4

Left
ventricular
hypertrophy

5

Bundle
branch block

6

Differential
diagnosis of
other atrial
arrhythmias

3. Laboratory tests

- Routine blood tests should be carried out at least once in patients with AF
- **Important parameters to assess include:**
 - Thyroid function (TSH)
 - Renal function
 - Hepatic function
 - Serum electrolytes & Complete blood count

4- Transthoracic echocardiography (TTE)


- **Non-invasive**, Used to identify:
 - **Size and functioning of atria and ventricles**
 - Ventricle hypertrophy
 - Pericardial disease
 - Valvular heart disease

5- Chest radiography

- When clinical findings suggest an abnormality chest radiography may be used to:
 - Evaluate pulmonary pathology and vasculature
 - Detect congestive heart failure
 - Assess enlargement of the cardiac chambers



★ 6- Holter monitor

- **Portable ECG device** (Now it can be done by modern wearable devices e.g. Smart watches)
- Useful for **diagnosis**¹ and **follow up** to see if the **treatment is effective**.²
- Continuous monitoring for a short period of time (typically 24-48 h)
- Useful for Detecting:
 - asymptomatic AF
 - Evaluating patients with paroxysmal AF
 - Associating symptoms with heart rhythm disturbance
 - Assessing response to treatment



★ 7- Transesophageal echocardiography (TEE)

- Ultrasound transducer positioned close to the heart using an endoscope-like device
- High quality images of cardiac structure and function:
 - Particularly the **left atrial appendage**, the **most common site of thrombi in patients with AF** (This is the only reliable method to rule out left atrial appendage thrombus)
- Not routinely used but **useful for**:
 - **Accurate assessment of risk of stroke**
 - Detection of low flow velocity ('smoke' effect)
 - Sensitive detection of atrial thrombi




1-e.g. If a patient presents to you saying that he has palpitations, but at the time of presentation he was in sinus rhythm, palpitations could be caused by anything so you give him this device and ask him to come back after 24hrs (sometimes 48 or 72hrs, depends on the episode) this device will monitor the ECG of the patient will help you determine whether the patient has AF or not.

2- e.g. You diagnosed a patient with AF and started treatment to control his symptoms, yet when he goes to the community he still complains of palpitations, so you give this device to monitor him and check if the AF is controlled or not and whether the patient requires a higher dose of treatment or not.

AF evaluation summary (From Dr slides)

[Click here for a very nice summary](#)



AF complications

Complication	General Info
Death	AF increases mortality 1.5 - 3.5 folds , due to sudden death , HF, comorbidities or stroke.
Stroke	20-30% of all ischemic strokes (preventable) and 10% of cryptogenic strokes are due to AF. Mainly due to cardioembolic or related to comorbid vascular atheroma.
Hospitalizations	10-40% of AF are hospitalized every year. For AF management, related to HF or MI or AF related symptoms also for treatment of associated complications.
Quality of life	More than 60% of patients. It's related to AF burden, comorbidities, psychological functioning and medication, distressed personality type.
Left ventricular dysfunction (LVD) & HF¹	LVD is found in 20-30% of all AF patients. Due to excessive ventricular rate, irregular ventricular contractions, a primary underlying cause of AF.
Cognitive decline and vascular dementia	HR 1.4/1.6 (irrespective of stroke history). Due to Brain matter lesions, inflammation or hypoperfusion and micro-embolism.
Depression	In 16-20% of patients (even suicidal ideation). Due to severe symptoms, decreased QoL and drug side effects.

1- How does AF cause HF? By decreasing ventricular filling, pulmonary congestion and dilation of ventricles.

AF Management

◀ Treatment of AF

- First treat the underlying cause then to initiate treatment for the arrhythmia you have to determine if the patient is hemodynamically stable or not.
 - How?** If there's **chest pain, shortness of breath, altered mental status (confusion)**, or a **systolic BP < 90**, then the patient is considered **unstable**. If they're unstable use electricity. If instead the patient has symptoms, but not any one of those listed above, the patient is **stable**. A patient who is stable has time to fix the rhythm. They're not going to die at this moment; pharmacotherapy can be used.

Hemodynamic stability

ABCD

Stable patients

- 1) Rate control (**B**B or **C**a++ blockers or **D**igoxin)
- 2) Rhythm control (**C**ardioversion)
- 3) **A**nticoagulation

Unstable patients

- 1) Do synchronized **electrical Cardioversion (Shock)** immediately.

◀ Pillars of AF management :

Pillars of AF Management



01

Risk factors management

control the risk factor that mention previously if the patient have any, for example; Alcohol, HTN, Obesity and Obstructive sleep apnea


02

Anticoagulation (Stroke prevention)

- Why we give anticoagulant in AF ? In AF the blood doesn't move → forming clot most commonly in the left atrial appendage → risk of stroke .
- We decide who to be anticipated and who's not through the **score system** (the more factor the patient has the more risk of stroke) :

1	CHA ₂ DS ₂ -VASc	Score
	C ongestive heart failure/LV dysfunction	1
	H ypertension	1
	A ge ≥ 75 years	2
	D iabetes mellitus	1
	S troke/TIA/TE	2
	V ascular disease [prior MI, PAD, or aortic plaque]	1
	A ge 65-74 years	1
	S ex category (female)	1

2	CHADS ₂ score	Patients (n = 1733)	Adjusted stroke rate (%/year) ^a (95% confidence interval)
	0	120	1.9 (1.2-3.0)
	1	463	2.8 (2.0-3.8)
	2	523	4.0 (3.1-5.1)
	3	337	5.9 (4.6-7.3)
	4	220	8.5 (6.3-11.1)
	5	65	12.5 (8.2-17.5)
	6	5	18.2 (10.5-27.4)



- Anticoagulant agents (dr : not gonna ask about in exam, just for your knowledge)
 - Warfarin > in rheumatic mitral stenosis and mechanical valves
 - New agents > any patient is treated with these agents except rheumatic MS and mechanical valve

438 slides

- Patient with **AF related Moderate or severe mitral stenosis or the presence of a mechanical heart valve should always be anticoagulated (With Warfarin)** because the risk is so high. In other patients, **CHA₂DS₂-VASc score** is used to assess the risk of stroke.
- **New Oral AntiCoagulants (NOAC) “First line treatment”**:¹
 - Dabigatran, Apixaban, Rivaroxaban, Edoxaban
- **Warfarin² (Coumadin®) “used in cases of Mitral stenosis or if pt has metal valve”**
- WATCHMAN® device or surgery: For removal/isolation of left atrial appendage.

1- The anticoagulants used for AF decrease the risk of stroke by almost 70% (except aspirin by 20%). In general the NOACs are better than warfarin why? Because of no need of monitoring (in warfarin you have to keep the INR between 1-3), No Drug food interaction (in warfarin, the patient can't eat Vit K containing food), Minimum Drug-Drug interaction. **BUT REMEMBER YOU SHOULDN'T GIVE THESE DRUGS TO PATIENTS WITH MITRAL STENOSIS OR MECHANICAL VALVE!! (The first line for them is VKA “warfarin”)**
 2- Most patients should be anticoagulated (INR 2.0-3.0) long term: the exception being young patients (<65 years) with lone AF. This latter group is treated with aspirin alone.

AF Management cont'

pillars of AF management cont.

03

Rate control (We aren't going to put the pt back to sinus rhythm we just want to make sure that the pt isn't tachycardic)

- Target heart rate for AF patients <110 bpm during activity and 60-80 bpm during rest .
- **Medication** (AV node blockers) : (BB and CCB have equal efficacy, choice depends on the pt situation, digoxin has lowest efficacy)
 - **Beta - blocker** Atenolol, Metoprolol (Used in cases of Graves disease, CHF, IHD, HTN)
 - **Calcium channel blocker (CCB)** Verapamil , diltiazem (Used if patient is Asthmatic)
 - **digoxin** (Used if patient is hypotensive)
- **procedure:**
 - **Pace and ablate** ¹> we only use this if we can't control the rate with medication .

04

Rhythm control convert the AF to the sinus rhythm

- **Medication (antiarrhythmic) : (PNR ² or maintenance)** [Click here](#) for a summary of all antiarrhythmics (EXTRA!!)
 - **Amiodarone**
 - **Sotalol**
 - **Flecainide**
- **procedure:**
 - **AF ablation**

438 slides

- Electrical Direct current (DC) cardioversion (More effective) or pharmacological cardioversion may be used.
 - Likelihood of successful cardioversion decreases with the duration of AF:
 - Pharmacological cardioversion is most effective when initiated within 7 days of AF onset.
 - It's advocated for younger, symptomatic and physically active patients.
 - **Pharmacologic cardioversion use depends on the presence or absence of underlying heart disease:** (It's mainly used if electrical cardioversion fails or not feasible)
 - A. Absent:** Class IC antiarrhythmics e.g. **IV Flecainide**, Propafenone Or Class IA e.g. Procainamide
 - B. Present:** Class III antiarrhythmics e.g. **IV Amiodarone**³ Or Dronedarone (alternative)
 - Others: Ablation⁴, Surgery (MAZE)
 - **AF for <48hr** → **Cardiovert immediately** either pharmacologically or **electrically** (or both) **then anticoagulate for 4 wks**
 - **AF for ≥48hr (or unknown)** → Either do **TEE** to check atrial thrombus **then cardiovert** (If no thrombus) **Or Give Anticoagulants for ≥3wks** → **Cardiovert** → **Give Anticoagulants again for at least 4wks.**
- Why do we give anticoagulants before cardioversion?** Because LAA thrombus can dislodge and cause stroke
Why do we give anticoagulants after cardioversion? Because cardioversion may lead to atrial stunning → Blood stasis → Thrombus formation.

3 important principles :

- Rate control is reasonable if asymptomatic (keep the patient on atrial fibrillation)
- Consider rhythm control for symptoms Rx (don't leave them on AF)
- Catheter ablation is a reasonable **first line** (We can just do it upfront without medical therapy)

1. In exceptional cases, poorly controlled and symptomatic AF can be treated by implanting a permanent pacemaker and then deliberately inducing complete AV nodal block with catheter ablation. This is known as the 'pace and ablate' strategy. **Pace and ablate** simply is procedure in which we take a catheter all the way up to the heart then burn the **AV- node!** burning the AV- node will lead to complete AV block (no more impulses will be conducted from the atrium to the ventricles), the goal is to protect the ventricle from the high atrial rhythm, after that we implant pacemaker to establish normal Rhythm in the ventricle
2. prescription is taken as needed, e.g. if the patient has episode of AF every 6 month > we don't want to let the patient take the drug daily so we tell him take it when you have AF .
3. The most effective antiarrhythmic drug for AF is amiodarone followed by flecainide and propafenone. The problem with amiodarone is its side effects (e.g. Photosensitivity skin discoloration, corneal deposits, thyroid dysfunction, alveolitis, nausea and vomiting, hepatotoxicity, peripheral neuropathy, torsade de pointes; potentiates digoxin and warfarin) so it's usually reserved for old people (>70), but for young people flecainide and propafenone are better.
4. Catheter ablation is sometimes used to restore and maintain sinus rhythm in resistant cases


CHA₂DS₂-VASc scoring system

- A scoring system used to identify which patient is at high risk of thromboembolic complications and will benefit from anticoagulation therapy.




16.23 CHA ₂ DS ₂ -VASc stroke risk scoring system for non-valvular atrial fibrillation		
Parameter		Score
C	Congestive heart failure	1 point
H	Hypertension history	1 point
A₂	Age ≥ 75 years	2 points
D	Diabetes mellitus	1 point
S₂	Previous stroke or transient ischemic attack (TIA)	2 points
V	Vascular disease	1 point
A	Age 65–74 years	1 point
Sc	Sex category female	1 point
Maximum total score		9 points

Annual stroke risk
 0 points = 0% (no prophylaxis required)
 1 point = 1.3% (oral anticoagulant recommended in males only)
 2+ points = > 2.2% (oral anticoagulant recommended)

0 → Low risk (No need for anticoagulants)
 1 → Moderate risk (the need of anticoagulants is controversial)
 2 or more → High risk (Requires anticoagulation)







AF Management Summary



No need to know this figure

Other Arrhythmias

Type of arrhythmia	General info	ECG	Management
<p>Atrial flutter الرفرفة الأذينية</p>	<ul style="list-style-type: none"> One irritable automaticity focus (macrocircuit) in the atria fires at about 250 to 350 bpm (typically very close to 300 bpm), giving rise to regular atrial contractions. Causes: Heart disease: Heart failure (most common association), rheumatic heart disease, CAD, COPD, Atrial septal defect (ASD) 	<p>ECG provides a saw-tooth baseline, with a QRS complex appearing after every second or third "tooth" (F wave).</p> 	<ul style="list-style-type: none"> Similar to Atrial fibrillation treatment except that we not manage the risk factors Most cases of flutter can be cured with radiofrequency catheter ablation of the re-entry circuit. Catheter ablation is first line¹
<p>Supraventricular tachycardia (SVT) تسرع القلب فوق البطيني (Benign)</p>	<ul style="list-style-type: none"> common , it occur in young patient without heart disease These are usually regular narrow-complex tachycardias and are characterised by a re-entry circuit or automatic focus involving the atria. The three types are: <ol style="list-style-type: none"> Atrioventricular <u>nodal</u> re-entrant tachycardia (AVNRT) (60%) when the focus is in the AV-node itself. Atrioventricular re-entrant tachycardia (AVRT) (30%) e.g. WPW, when there is an accessory pathway. Atrial tachycardia (10%) if the focus is in the atrium). 	<ul style="list-style-type: none"> in general > regular narrow-complex tachycardias. 1. AVNRT⁴³⁸: QRS complexes are narrow and the P waves cannot be seen.  <ul style="list-style-type: none"> 2. AVRT⁴³⁸ : The tachycardia P waves (arrowed) are clearly seen after narrow QRS complexes 	<p>Acute :</p> <ul style="list-style-type: none"> - Carotid massage - valsalva manoeuvre - Medical (IV adenosine, CCB, BB, flecainide) <p>Severe haemodynamic compromise : DC cardioversion</p> <p>Chronic :</p> <ul style="list-style-type: none"> - catheter ablation² - Medical therapy - watchful waiting³ <p>Prophylaxis : BB, CCB, flecainide</p>
<p>Ventricular fibrillation (VF) الرجفان البطيني</p>	<ul style="list-style-type: none"> Multiple foci in the ventricles fire rapidly, leading to a chaotic quivering of the ventricles and no cardiac output. Most episodes of VFib begin with VT (except in the setting of acute ischemia/ infarction). there are 3 main causes : <ul style="list-style-type: none"> o ischemia o cardiomyopathy o primary arrhythmia⁴ 	<ul style="list-style-type: none"> No atrial P waves can be identified. No QRS complexes can be identified. In sum, no waves can be identified; there is a very irregular rhythm. 	<ul style="list-style-type: none"> - VF is a medical emergency! Immediate defibrillation and CPR are indicated, Fatal if untreated. - DC cardioversion immediately. If the equipment isn't ready, start CPR until it is. - If Persistent: Epinephrine - Refractory: IV amiodarone followed by shock



1- The procedure is much easier than in AF > so almost all patient with atrial flutter will undergo this procedure

2- Because it appears in young healthy persons, we tend to do ablation so they don't suffer with medication for the rest of their life

3- If the patient doesn't have frequent episodes > nothing bad will happen other than the symptoms

4- For example; Long QTs syndrome (LQTS), and Brugada syndrome


Other Arrhythmias

Type of arrhythmia	General info	ECG	Management
Ventricular tachycardia (VT) تسرع القلب البطيني	<ul style="list-style-type: none"> Defined as rapid and repetitive firing of three or more PVCs in a row, at a rate of between 100 and 250 bpm CAD with prior MI is the most common cause has two type : <ul style="list-style-type: none"> VT with structural heart disease (e.g. ischemic cardiomyopathy, hypertrophic cardiomyopathy)¹ VT without structural heart disease (i.e. idiopathic ventricular tachycardia) 	<ul style="list-style-type: none"> wide and bizarre QRS-complex, regular tachycardia 	<ul style="list-style-type: none"> Treat the underlying cause. Automatic implantable defibrillators <ul style="list-style-type: none"> VT with structural heart disease : <ul style="list-style-type: none"> ICD (device implanted under skin and there is a wire goes to the heart and detect the fast rhythm then correct it by shocking the patient) it does not treat VT it just prevents it) medication (BB , antiarrhythmic (AA)) Ablation² VT without structural heart disease : <ul style="list-style-type: none"> medical therapy (bb +/- AA) Ablation
Wolff-Parkinson-White Syndrome (WPW)	<ul style="list-style-type: none"> In AF the atria can reach 600 bpm while the ventricles are kept at a very lower rate due to AV nodal delay, in WPW, the atria beats and the impulses travel down the ventricles through accessory pathway (no block of impulse). Therefore if a patient is known to have WPW and develops AF he will die due to sudden cardiac death secondary to VFib Wolff-Parkinson-White (WPW) syndrome patients who present in AF with rapid ventricular response: If baseline ECG shows a delta wave or if the current ECG shows wide, bizarre QRS complexes during AF, avoid AV nodal blocking agents (β-blockers, CCBs, adenosine, digoxin). 	<ul style="list-style-type: none"> Delta-wave (the p-wave is short and it looks like it is fused with QRS-complex) The delta-wave appears due to the accessory pathway (this accessory pathway is faster than AV-node) > that's why PR-interval is short 	<ul style="list-style-type: none"> If the accessory pathway is capable to conduct fast impulses then there will be a risk of Sudden cardiac death therefore we do cath ablation to this accessory pathway Even if you find it by chance after doing ECG for any reason you should assess the accessory pathway and don't ignore it because it's asymptomatic . How to know if the accessory pathway is capable or not : <ol style="list-style-type: none"> Run on treadmill : increase the heart rate if the delta wave disappears that means it is not capable . Holter monitor : and see when to lose the delta wave The treatment of choice is IV procainamide or ibutilide, which slows conduction in the entire atrium. If AV nodal blocking agents are given in this situation, the atrial impulses in rapid AF can proceed down the accessory pathway and cause VF and death.

[Click here](#) for more info


1- They are at risk of Sudden cardiac death .

2- Decision is complicated, for example if the patient has ischemic cardiomyopathy> ablation is good and successful most of the time, but in case of hypertrophic cardiomyopathy the muscle is thick and Unsuccessful rate is higher .



1. Check if the patient has dyspnea, chest pain, confusion or systolic BP <90 (hypotension)
2. First step in treating a stable patient with arrhythmia is to determine what type of arrhythmia the patient has by **ECG**.
3. To make sure that patient does not have left atrial thrombus LAT (through Echocardiography(TEE)); because if he/she does have clot and you shock him/her it will lead to a massive stroke
4. Why? To increase the vagal tone which will block the AV-node > and it will be enough to break the circuit > terminate SVT
5. AV-blocker that works for few seconds which will break the circuit and terminate SVT
6. **Most of the time we can terminate SVT without the use of cardioversion through Carotid massage or Adenosine so you must consider them first b/c they are non-invasive and very effective methods.**

A 75-year-old man with a past medical history of diabetes mellitus and hypertension presents to the emergency department complaining of the sudden onset of shortness of breath accompanied by palpitations beginning 6 hours ago. The palpitations last for approximately 10 minutes at a time and recur at least once an hour. He admits to a 5-year history of intermittent similar symptoms. He denies chest pain, cough, or light-headedness. He takes no medications and does not smoke cigarettes, but admits to drinking six to eight beers per day over the past 3 days. Vital signs include a temperature of 37.2°C (98.9°F), blood pressure of 135/90 mm Hg, pulse rate of 130/min, and respiratory rate of 22/min. The patient is speaking in full sentences, has a midline trachea, and has no inspiratory rales, dullness to percussion, or increased tactile fremitus over the lung fields. His heart examination is notable for an irregularly irregular rhythm without murmurs; there is no chest wall tenderness. An ECG is shown in Figure



Q1: What's the most likely diagnosis?

Atrial fibrillation (AF). This is the only common arrhythmia in which the ventricular rate can be **rapid and irregular**. Atrial flutter is often confused with AF; however, atrial flutter has a rapid regular ventricular response with a rate about 150 beats per minute and the characteristic ECG finding of flutter waves.

Q2: How is this condition classified?

- Paroxysmal
- Persistent
- Long-standing Persistent
- Permanent
- Lone
- Non-valvular

Q3: What are the possible causes?

Hypertensive heart disease, HF and 3 PIRATES


Q4: Describe his ECG.

- Absent P waves, only a fine oscillation of the baseline (so-called fibrillation or f waves)
- Fast and irregular ventricular response (QRS complex) → irregular RR intervals (irregularly irregular rhythm)
- Increased heart rate (Tachyarrhythmias)

Q5: What's the appropriate treatment for this condition in general?

ABCD

- **Unstable:** In patients with acute-onset AF and hemodynamic instability, electrical cardioversion is indicated.
- **Stable:** If the duration of the AF is unknown or > 48 hours, or the patient is at high risk of embolization, transesophageal echocardiography (TEE) is performed to locate atrial clots prior to cardioversion. Alternatively, cardioversion may be delayed for 3 weeks for anticoagulation to minimize the risk of embolization in a stable patient then followed by another 4 weeks of anticoagulation.



Setting: ED

CC: "I feel a fluttering in my chest"

VS: RR: 32 breaths/minute: BP: 118/88 mmHg: P: 128bpm, irregularly irregular; T: 98.1°F

HPI: A 42-year-old gastroenterologist comes to the office with 1 single day of palpitations and fluttering in his chest. He has never had this before. He denies chest pain, lightheadedness, or shortness of breath. He is very anxious. He drinks a large amount of vodka at night four to five times a week and frequently travels to present papers at international meetings.

PMHx: None


Medications: None

PE:

- Neuro: Normal
- Cardio: No murmurs, rubs or gallops
- Abdomen: normal
- Extremities: no edema

Initial orders:

- ECG
- CHEM-7
- CXR
- Oximeter
- CBC



As you move the clock forward 10-20 min, all the laboratory results come back as normal except for the ECG, which shows AF at a rate of 125 to 130 bpm.

Q1: When is immediate electrical cardioversion for AF correct?

Electrical cardioversion is indicated for tachyarrhythmias when there is life-threatening hemodynamic instability, such as chest pain, shortness of breath, decreased systolic BP or hemodynamically related altered mental status (confusion). Although uncomfortable, none of other symptoms described in the case here are severe enough to put the patient through the risk and discomfort of electrical cardioversion. Pallor, anxiety, sweating and palpitations are all subjective, hard to measure and not life-threatening.

Q2: What's the first line treatment for this case?

Rate control with either a beta-blocker, CCB or digoxin is the first step in managing rapid AF. Rate control is more important than trying medications such as antiarrhythmics to chemically convert AF into normal sinus rhythm. Anticoagulation is not needed for AF present for <48hrs. The echocardiogram results may help tell who needs anticoagulation, but controlling the rapid ventricular response is not dependent on echocardiogram findings.

After metoprolol or diltiazem, the patient's HR reduces to 80bpm. All symptoms resolve. admittance to the ICU is not necessary and the patient is placed on a regular hospital ward. Heparin is not necessary. The echocardiogram shows a normal heart size and shape with no significant valvular disease and no thrombi. The AF does not stop, but the rate remains controlled and there are no symptoms.

Q3: What's the best therapy?

- A- Warfarin
- B- Dabigatran
- C- Aspirin
- D- Rivaroxaban

A: C. This patient has a CHADS score of 0 so aspirin alone is sufficient. If CHADS score is 0 or 1, Aspirin alone is enough, if >2 give NOAC or Warfarin.

Summary

Atrial fibrillation

General info

- Prevalence **rise with age**. It's also **more common in males**.

Pathogenesis

- Multiple interacting re-entry microcircuits looping around the atria**, most often located within pulmonary veins

Causes

Hypertensive heart disease, HF and 3 PIRATES

AF rhythm & ECG changes

- Absent P waves**, only a fine oscillation of the baseline (so-called fibrillation or f waves)
- Fast and irregularly irregular ventricular response → irregular RR intervals
- Increased heart rate (Tachyarrhythmias)
- irregularly irregular rhythm.**

AF begets AF

Electrical remodeling

Electrophysiological changes occur in the atria within a few hours of the onset of AF, that tend to maintain fibrillation

Structural remodeling

When AF persists for a period of months, structural remodeling occurs with **atrial fibrosis and dilation** that further predispose to AF

S & S

- Irregularly irregular pulse**
- palpitation, breathlessness and fatigue
- Thromboembolic TIA → **Stroke**

Classification

- Paroxysmal AF** → AF <7d
- Persistent AF** → >AF >7d
- Long-standing persistent AF** → >AF 12 mo
- Permanent AF** → Joint decision between pt and clinician not to pursue rhythm control
- Lone AF** → AF without clinical/ECG evidence of cardiopulmonary disease
- Non valvular AF**

Diagnosis

- History & Physical examination:** Clinical type of AF, presence of underlying heart diseases, Response to any treatments administered, Triggers, family history.
- ECG:** Abnormal Rhythm (verify AF), LVH, pre-excitation, Prior MI, Bundle branch block
- Transthoracic echocardiogram (TTE):** Size and functioning of atria and ventricles, LV hypertrophy, VHD
- Blood tests** to check the thyroid, renal and hepatic
- Holter monitor:** If diagnosis of type of arrhythmia is in question.
- Chest X-ray**
- Transesophageal echocardiogram (TEE):** To identify LA thrombus in LAA, to guide cardioversion

Complications

- Stroke**, Death, Hospitalizations
- Impaired quality of life, Left ventricular dysfunction (LVD), Cognitive decline and vascular dementia

Management

Stable

Unstable

- Rate control:** **B**B or **C**a⁺⁺ blockers or **D**igoxin
- Rhythm control:**
 - AF for <48hr → **C**ardiovert immediately either pharmacologically or electrically (or both)
 - AF for >48hr → Either do TEE to locate atrial thrombus then cardiovert Or Give Anticoagulants for 3wks → **C**ardiovert → Give Anticoagulants again for 4wks.
- A**nticoagulation: NOAC or Warfarin

Synchronized **electrical Cardioversion (Shock)** immediately

ABCD

Lecture Quiz

Q1: A third-year medical student has been reading about the dangers of excessive anticoagulation and bleeding potential. He reviews the charts of several patients with atrial fibrillation currently taking Coumadin. Which of the following patients is best suited to have anticoagulation discontinued?

- A- A 45-year-old man who has normal echocardiographic findings and no history of heart disease or hypertension, but a family history of hyperlipidemia
- B- A 62-year-old man with mild chronic hypertension and dilated left atrium, but normal ejection fraction
- C- A 75-year-old woman who is in good health except for a prior stroke, from which she has recovered nearly all function
- D- A 52-year-old man with orthopnea and paroxysmal nocturnal dyspnea

Q2: A 48-year-old woman is noted to have atrial fibrillation with a ventricular heart rate of 140 bpm. She is feeling dizzy and dyspneic with a systolic blood pressure of 75/48 mm Hg. Which of the following is the most appropriate next step?

- A- Intravenous digoxin
- B- DC cardioversion
- C- Vagal maneuvers
- D- Intravenous diltiazem (Cardizem)

Q3: A 45-year-old woman is noted to have dizziness, pounding of the chest, and fatigue of 3 hours' duration. On examination, she is noted to have a blood pressure (BP) of 110/70 mm Hg and heart rate of 180 bpm. She is noted on ECG to have atrial fibrillation, and a prior baseline ECG showed delta waves. The ER physician counsels the patient regarding cardioversion, but the patient declines. Which of the following is the best therapy for her condition?

- A- Digoxin
- B- Angiotensin-converting enzyme (ACE) inhibitor
- C- Calcium channel blocker
- D- Procainamide

Q4: A 68-year-old man presents with a 2-week history of increasing dyspnea and is found to be in atrial fibrillation with a ventricular rate of 120 beats per minute. The most appropriate test to evaluate for left atrial thrombus before cardioversion is

- A- ECG
- B- Transthoracic echocardiography
- C- No testing is needed
- D- Transesophageal echocardiography

Q5: A 79-year-old woman is admitted to the coronary care unit (CCU) with unstable angina. She is started on appropriate medication to reduce her cardiac risk. She is hypertensive, fasting glucose is normal and cholesterol is 5.2. She is found to be in atrial fibrillation. What is the most appropriate treatment?

- A- Aspirin and clopidogrel
- B- Digoxin
- C- Warfarin
- D- Cardioversion

Q6: A 62-year-old male presents with palpitations, which are shown on ECG to be atrial fibrillation with a ventricular rate of approximately 130/minute. He has mild central chest discomfort but is not acutely distressed. He first noticed these about 3 hours before coming to hospital. As far as is known this is his first episode of this kind. Which of the following would you prefer as first-line therapy?

- A- Anticoagulate with heparin and start digoxin at standard daily dose
- B- Attempt DC cardioversion
- C- Administer bisoprolol and verapamil, and give warfarin
- D- Attempt cardioversion with IV flecainide

GOOD LUCK!

*This work was originally done by **438 Medicine team:***

Team Leaders

- Raghad AlKhashan - Mashal AbaAlkhail
- Amirah Aldakhilallah - Ibrahim AlAsous

Member : Masha AbaAlkhail

Note taker : Raghad Alkhashan

*Edited by **439 Medicine team:***

Team Leaders

- Shaden Alobaid - Hamad Almousa
- Ghada Alabdi - Naif Alsulais

Member : Nouf Alsubaie

Note taker : Abdullah Alanzan



CONTACT US THROUGH OUR EMAIL :

MEDICINE439@GMAIL.COM