



Measures of Disease Frequency, Impact and Effects

Dr. Shatha Alduraywish, MBBS; MEpi; PhD

Assistant Professor, Epidemiologist

Department of Family and Community Medicine

College of Medicine, KCUH

King Saud University

Salduraywish@ksu.edu.sa

Mar 2020


Learning Objectives

By the end of this session, the students will be able to:

- Understand the concept of the common epidemiological measures of disease frequency and effect.
- Identify the appropriate use of these measures
- Interpret the measures of disease frequency, impact and effect


Table 1.1. Questions relevant for epidemiological enquiry

Disease definition	What characteristics or combination of characteristics best discriminate disease from non-disease?
Disease occurrence	What is the rate of development of new cases in a population? What is the proportion of current disease within a population? What are the influences of age, sex, time and geography on the above?
Disease causation	What are the risk factors for disease development and what are their relative strengths with respect to an individual and a population?
Disease outcome	What is the outcome following disease onset and what are the risk factors, including their relative strengths, for a poor outcome?
Disease management	What is the relative effectiveness of proposed therapeutic interventions? (Included within this are health service research questions related to the relative effectiveness of proposed health service delivery options.)
Disease prevention	What is the relative effectiveness of proposed preventive strategies including screening?



Measures for Disease Occurrence

- **Proportions:**
 - Prevalence
 - Incidence proportion
- **Rates:**
 - Incidence rates
- **Ratio:**
 - odds for a certain disease



Proportion, Rate and Ratio

▪ Proportions

- They are dimensionless (**do not have a unit** of measure, because the unit of measure in the denominator is the same as the numerator)
- Always lies between 0 and 1


▪ Rates

- Denominator is measured in **time units**
- Can exceed 1 if no. of new cases > person-time spent at risk

▪ Ratio


- **Compares** between two measures (two rates, odds or proportions)
- What is counted in numerator **isn't always** in the denominator

Measures of Disease Frequency



Prevalence

- The amount of a disease in a population at a given point in time.



Point Prevalence

The proportion of the **population** that has the disease at a **specific point** in time

Point Prevalence

$$= \frac{\text{Number of **current** cases at a specific point in time}}{\text{Total population at that same point in time}}$$

“**Current cases**” means new and pre-existing cases (all the cases that were there at that point in time)

Period Prevalence

- The proportion of the **population** that has the disease during **a specified period** of time

Period Prevalence

$$= \frac{\text{Number of **current** cases during a specific period of time}}{\text{Average or mid-interval population}}$$

Prevalence (Example)

	Asthma	No Asthma	Total
Non-smokers	40	360	400
smokers	30	170	200
Total	70	530	600

What is the prevalence of asthma among
(1) Smokers and
(2) Non-smokers?

Prevalence of asthma among smokers = $30/200 = 15\%$

Prevalence of asthma among non-smokers = $40/400 = 10\%$

Incidence Proportion


Incidence Proportion =

Number of new cases

total population at risk at the beginning of the study

- **The population at risk** is a well-defined population that is **free** of the disease at the beginning of the study and has certain characteristics that put them at risk for developing the disease

Follow-up in Study



of people at risk at baseline?

of cases developed during the 6 year follow-up period?

Total person-time at risk?

Incidence Rate

Incidence Rate=

*Number of **new cases***

*the total **person time** at risk over the study period of time*

- Here we are taking into consideration the **time** that each person spent being at risk before developing the disease
- By contrast the incidence proportion only considers the total population at risk without also incorporating time in the equation

Rate vs. Risk

A study followed **3,000** males ages 45 years and older for **5 years** to assess the development of MI.

During the study period, **150** men developed MI, who accumulated a total person-time of **14,625 person-years**.

What is the *incidence proportion* after 5 years?

What is the *incidence rate* after 5 years (rate)?

Prevalence vs. Incidence

- Prevalence → Cross-sectional study (Survey)
- Incidence → Cohort study

Comparisons between Prevalence and Incidence

Prevalence

- One point in time; easy to measure
- Proportion or %
- Numerator: count of people with disease
- Denominator: count of total population at risk
- No time component

Incidence

- Involves time; difficult to measure
- Measured as either *rate* or *proportion*
- Numerator: count of people who develop disease during follow-up
- Denominator:
 - (prop.) People at risk and disease free
 - (rate) Person-time at risk.

Measures of Effect (Associations)



Measures of Effect (Associations)

- Risk
- Odds
- Risk Ratio
- Odds Ratio
- Relative Risk Reduction and Absolute Risk Reduction
- Number needed to Treat and Number needed to Harm

2X2 Table

		Outcome		Total
		Yes	No	
Exposure	Yes	a	b	a + b
	No	c	d	c + d
Total		a + c	b + d	N

Odds

- The ratio of the probability of occurrence of an event to that of non-occurrence
- Odds in Exposed = a/b
- Odds in unexposed, “Baseline odds” = c/d

		Outcome		Total
		Yes	No	
Exposure	Yes	a	b	a + b
	No	c	d	c + d
Total		a + c	b + d	N

Odds Ratio (OR)

	Outcome	Outcome	Total
Exposure	Yes	No	
Yes	a	b	a + b
No	c	d	c + d
Total	a + c	b + d	N

$$\text{Odds ratio} = \frac{a/b}{c/d} = \frac{ad}{cb}$$

Odds Ratio

	Ear Infection	Ear Infection	Total
Swimming	Yes	No	
Yes	40	60	100
No	5	95	100
Total	45	155	200

Odds of getting infection for **swimmers**: $40/60 = 0.67$

Odds of getting infection for **non-swimmers**: $5/95 = 0.052$

$$\text{Odds ratio} = \frac{0.67}{0.052} = 12.7$$

Risk

- Probability that an event will occur
 - Risk in Exposed = $a / (a+b)$
 - Risk in unexposed, “Baseline risk” = $c / (c+d)$

		Outcome		Total
		Yes	No	
Exposure	Yes	a	b	a + b
	No	c	d	c + d
Total		a + c	b + d	N

Relative Risk (RR)

		Outcome		Total
		Yes	No	
Exposure	Yes	a	b	a + b
	No	c	d	c + d
Total		a + c	b + d	N

$$\text{Relative Risk} = \frac{a/(a+b)}{c/(c+d)}$$

Relative Risk

Interpretation:

How many **times** more likely it is that someone who is exposed to something will develop a certain disease compared to someone who is not exposed.

- 1 → no difference between the groups
- <1 → reduced the risk (protective)
- >1 → increase the risk

Relative Risk

	Ear Infection	Ear Infection	Total
Swimming	Yes	No	
Yes	40	60	100
No	5	95	100
Total	45	155	200

Risk of getting infection **for swimmers**: $40/100 = 0.4$

Risk of getting infection for **non-swimmers**: $5/100 = 0.05$

$$\text{Relative Risk} = \frac{0.4}{0.05} = 8$$

Relative Risk

RR doesn't tell you the **magnitude of benefit of treatment.**

It only tells there is increase or decrease risk in experiment group compared to control group.

Absolute Risk Reduction (ARR)

- Risk Difference.
- $ARR = RR \text{ (exposed)} - RR \text{ (Unexposed)}$
- It tells the **magnitude of benefit**

Interpretation

if 100 patients were treated with experiment treatment, (x) cases of outcome can be prevented.

Example: if **ARR = 15%** in comparing ACEI vs placebo in decreasing IHD. This means if **100 patients were treated with ACEI, 15 cases of IHD can be prevented compared to placebo.**

- If ARR equals **0**, then there is no difference between experiment and control.

Relative Risk Reduction (RRR)

- $RRR = 1 - RR$
- It tells how much the experiment treatment is reducing the chance of having outcome in single treated patient.

Interpretation:

- Using experiment treatment will relatively reduce the risk of having the outcome by (%) compared to control treatment.

Example

if **RRR = 70%** in comparing ACEI vs placebo in decreasing IHD.

- This means **treatment with ACEI will relatively reduce the risk of having IHD by 70% compared to placebo**; i.e.
- in person using ACEI, his chance of having IHD will be reduced by 70%

Number Needed to Treat (NNT)

- Number of persons who would have to receive an intervention for 1 to benefit.
- $NNT = 1/ARR$

Measures of Impact



© Can Stock Photo - csp35840122

Measures of Impact

- Measures of association providing information about **absolute effects** of exposure
- Reflect apparent **contribution** of an exposure to the **frequency** of disease

Attributable Risk (AR)

- Quantifies disease burden in **exposed group** attributable to exposure
- **Provides answers to**
 - What is the risk attributed to the exposure?
 - What is the excess risk due to the exposure?
- Calculated as **risk difference (RD)**

Attributable Risk (Example)

Attributable Risk for Fast driving

	Dead	Not Dead	Total
Drive Fast	100	1900	2000
Drive Slow	80	7920	8000
Total	180	9820	10000

$$AR = 100/2000 - 80/8000 = 0.04 = 40\%$$

Summary

- Measures of Disease Frequency
 - Prevalence
 - Incidence
- Measures of Effect (Associations)
 - Risk
 - Odds
 - Risk Ratio
 - Odds Ratio
 - Relative Risk Reduction and Absolute Risk Reduction
 - Number needed to Treat
- Measures of Impact
 - Attributable Risk (AR)

A white, hand-drawn style thought bubble sticker is centered on a corkboard background. The sticker has a soft, irregular shape with a small tail pointing downwards. Inside the bubble, the words "Thank you!!" are written in a bold, black, handwritten-style font. The corkboard background is a warm, textured brown color.

Thank
you!!