

# Urologic Disorders

Abdulaziz Althunayan

Consultant Urologist

Assistant professor of Surgery



# Urologic Disorders

- Urinary tract infections
- Benign Prostatic Hyperplasia and voiding dysfunction
- Scrotal disorders

# Urinary tract infections

- Urethritis
- Epididymitis/orchitis
- Prostatitis
- cystitis
- Acute Pyelonephritis
- Chronic Pyelonephritis
- Renal Abscess

# URETHRITIS

## ■ S&S

- urethral discharge
- burning on urination
- Asymptomatic

## ■ **Gonococcal vs. Nongonococcal**

DX:

- incubation period(3-10 days vs. 1-5 wks)
- Urethral swab
- Serum: Chlamydia-specific ribosomal RNA

# URETHRITIS

Table 17-1. CLASSIC URETHRITIS

	Gonorrhea	Chlamydia
Organism	<i>Neisseria gonorrhoeae</i>	<i>Chlamydia trachomatis</i>
Organism type	Gram-negative diplococci	Intracellular facultative anaerobe
Incubation period	3-10 days	1-5 wk
Urethral discharge	Usually profuse, purulent	Usually scant
Asymptomatic carriers	40%-60%	40%-60%
Diagnostic test	Ligand chain reaction	Polymerase/ligand chain reaction
Other tests	Gram stain Culture	Culture Immunoassay
Recommended treatment	Ceftriaxone 125 mg IM once <i>plus</i> Azithromycin 1 g PO <i>or</i> Doxycycline 100 mg PO bid × 7 days	Azithromycin 1g PO <i>or</i> Doxycycline 100 mg PO bid × 7 days
Alternative treatment	Cefixime 400 mg PO <i>or</i> Ciprofloxacin 500 mg PO <i>or</i> Ofloxacin 400 mg PO <i>plus</i> Azithromycin 1 g PO <i>or</i> Doxycycline 100 mg PO bid × 7 days	Erythromycin 500 mg PO qid 7 days <i>or</i> Erythromycin ethylsuccinate 800 mg PO qid × 7 days <i>or</i> Ofloxacin 300 mg PO bid × 7 days

# Epididymitis

- Acute : pain, swelling, of the epididymis <6wk
- chronic :long-standing pain in the epididymis and testicle, usu. no swelling.
- DX
  - Epididymitis vs. Torsion
  - U/S
  - Testicular scan
  - Younger : *N. gonorrhoeae* or *C. trachomatis*
  - Older : *E. coli*

# Epididymitis

**Table 17-3. TREATMENT OF ACUTE EPIDIDYMO-ORCHITIS**

## **Epididymo-Orchitis Secondary to Bacteriuria**

1. Do urine culture and sensitivity studies
2. Promptly administer broad-spectrum antimicrobial agent (e.g., tobramycin, trimethoprim-sulfamethoxazole, quinolone antibiotic)
3. Prescribe bed rest and perform scrotal evaluation
4. Strongly consider hospitalization
5. Evaluate for underlying urinary tract disease

## **Epididymo-Orchitis Secondary to Sexually Transmitted Urethritis**

1. Do Gram stain of urethral smear
2. Administer ceftriaxone, 250 mg IM once; then tetracycline, 500 mg PO qid for at least 10 days, or doxycycline, 100 mg PO bid for at least 10 days
3. Prescribe bed rest and perform scrotal evaluation
4. Examine and treat sexual partners

Adapted from Berger RE: Urethritis and epididymitis. *Semin Urol* 1983;1:143.



# Prostatitis

- Syndrome that presents with inflammation± infection of the prostate gland including:
  - Dysuria, frequency
  - dysfunctional voiding
  - Perineal pain
  - Painful ejaculation



# Prostatitis

**Table 15–1. CLASSIFICATION SYSTEM FOR THE PROSTATITIS SYNDROMES**

Traditional	National Institutes of Health	Description
Acute bacterial prostatitis	Category I	Acute infection of the prostate gland
Chronic bacterial prostatitis	Category II	Chronic infection of the prostate gland
N/A	Category III chronic pelvic pain syndrome (CPPS)	Chronic genitourinary pain in the absence of uropathogenic bacteria localized to the prostate gland with standard methodology
Nonbacterial prostatitis	Category IIIA (inflammatory CPPS)	Significant number of white blood cells in expressed prostatic secretions, postprostatic massage urine sediment (VB3), or semen
Prostatodynia	Category IIIB (noninflammatory CPPS)	Insignificant number of white blood cells in expressed prostatic secretions, postprostatic massage urine sediment (VB3), or semen
N/A	Category IV asymptomatic inflammatory prostatitis (AIP)	White blood cells (and/or bacteria) in expressed prostatic secretions, postprostatic massage urine sediment (VB3), semen, or histologic specimens of prostate gland


N/A, not applicable.

# Prostatitis

## ■ Acute Bacterial Prostatitis :

- Rare
- Acute pain
- Storage and voiding urinary symptoms
- Fever, chills, malaise, N/V
- Perineal and suprapubic pain
- Tender swollen hot prostate.
- Rx : Abx and urinary drainage

# Chronic Prostatitis/Chronic Pelvic Pain Syndrome



# cystitis

## ■ S&S:

- **dysuria, frequency, urgency, voiding of small urine volumes,**
- **Suprapubic /lower abdominal pain**
- **± Hematuria**
- **DX:**
  - **dip-stick**
  - **urinalysis**
  - **Urine culture**

**Table 14–10. TREATMENT REGIMENS FOR ACUTE CYSTITIS**

Circumstances	Route	Drug	Dosage (mg)	Frequency per Dose	Duration (days)	
<b>Women</b>						
Healthy	Oral	Ciprofloxacin	500	Every 12 hr	3	
		Enoxacin	400	Every 12 hr		
		Levofloxacin	500	Every day		
		Lomefloxacin	400	Every day		
		TMP-SMX	160–800	Every 12 hr		
		TMP	100	Every 12 hr		
		Microcrystalline nitrofurantoin	100	Four times a day		
		Norfloxacin	400	Every 12 hr		
		TMP–SMX	160–800	Every 12 hr		7
		or	As above	As above		
Symptoms for >7 days, recent urinary tract infection, age >65 yr, diabetes, diaphragm use	Oral	Fluoroquinolone			7	
		Amoxicillin	250	Every 8 hr		
		Cephalexin	500	Four times a day		
		Microcrystalline nitrofurantoin	100	Four times a day		
Pregnancy	Oral	TMP-SMX	160–800	Every 12 hr	7	
<b>Men</b>						
Healthy and <50 years old	Oral	TMP-SMX	160–800	Every 12 hr	7	
		or				
		Fluoroquinolone	As above	As above		

TMP, trimethoprim; TMP-SMX, trimethoprim-sulfamethoxazole.

Modified from Stamm WE, Hooton TM: Management of urinary tract infections in adults. N Engl J Med 1993; 329: 1328–1334. Copyright 1993 Massachusetts Medical Society. All rights reserved.

# Pyelonephritis

- Inflammation of the kidney and renal pelvis
- S&S :
  - Chills
  - Fever
  - Costovertebral angle tenderness (flank Pain)
  - GI:abdo pain, N/V, and diarrhea
  - Gr-ve sepsis
  - Dysuria, frequency

# Pyelonephritis

## ■ Investigation:

- Urine C&S :+VE(80%)
  - *Enterobacteriaceae (E. coli), Enterococcus*
- Urinalysis:↑ WBCs, RBCs,Bacteria
- (±) ↑serum Creatinine
- CBC : Leukocytosis




# Pyelonephritis


## ■ Imaging:

- IVP
- U/S
- CT

Symptoms and Signs of Pyelonephritis  
(Fever, Flank Pain, Leukocytosis)



# Voiding Dysfunction



# LUTS

## STORAGE

(irritative)

- Dysuria
- Frequency
- Nocturia
- Urgency
- Incontinence

## VOIDING

(obstructive)

- Hesitancy
- Weak stream
- Straining
- Intermittency
- Dripping
- Retention

# Voiding Dysfunction


## ■ Failure to store

- Bladder problems
  - overactivity
  - Hypersensitivity
- Outlet problem
  - Stress incontinence
  - Sphincter deficiency
- combination

## ■ Failure to void

- Bladder problems
  - Neurologic
  - Myogenic
  - idiopathic
- Outlet problem
  - BPH
  - Urethral stricture
  - Sphincter dyssynergia
- combination

# Over Active Bladder



# Work up

- History
- Physical exam
- UA
- C/S
- US




# Treatment

- Behavioral
- Pelvic floor exercise
- Anti-cholinergics
- beta-3 agonist


# Benign Prostatic Hyperplasia BPH

## ■ Clinically:

- LUTS
- poor bladder emptying
- urinary retention
- urinary tract infection
- Hematuria
- Renal insufficiency



Normal Prostate




Enlarged Prostate

# Benign Prostatic Hyperplasia

## ■ Physical Examination


- 1-DRE 2- Focused neurologic exam
  - Prostate Ca
  - rectal Ca
  - anal tone
  - neurologic problems
- Abdomen: distended bladder



# Benign Prostatic Hyperplasia

- Urinalysis , culture
  - UTI
  - Hematuria
- Serum Creatinine
- Serum Prostate-Specific Antigen
- Flow rate
- Ultrasound (Kidney, Bladder And Prostate)

# US




# PVR





# Uroflowmetry





# Complications

- bladder stones
- UTI
- bladder decompensation
- incontinence
- upper tract deterioration
- hematuria
- AUR

# Benign Prostatic Hyperplasia

## ■ Treatment options

### – medical therapy

#### ■ $\alpha$ -Adrenergic Blockers

- Tamsulocin
- Alfuzocin
- Terazocin



#### ■ Androgen Suppression

- Finasteride
- Dutasteride



# Benign Prostatic Hyperplasia

## ■ Surgical Rx

- Endoscopic
- Transurethral Resection of the Prostate TURP
- Laser ablation
- prostatic stents




# Open Prostatectomy





# Hydrocele

- hydrocele occurs in males when fluid fills the scrotum.
- Fluid can surround one or both testicles, causing swelling in the scrotum.
- More common in baby boys, it may also occur in adult men.
- Imaging
  - Ultrasound
- Management
  - Hydrocelectomy






# Hydrocele sac






# Varicocele



- Most identified common cause of male infertility.
- It affects 20% of male population.
- Not all men with varicocele are infertile.
- It can be graded:
  1. Palpable with Valsalva maneuver.
  2. Palpable without Valsalva maneuver.
  3. Visible.

# Varicocele

- More common on the left side.
- It impairs fertility by increase intratesticular temperature.
- Treatment of varicocele indicated in:
  - Infertility with abnormal semen parameters.
  - Testicular pain
  - Impaired testicular volume.
- Treatment:
  - ligation (open, microscopic or lap)
  - Angioembolization



# varicoceles







# Varicoceles microligation

