

# Presentation &Management Of Common Thoracic Diseases


Dr. Waseem HAJJAR MD, FRCS, FCCP. Professor & Consultant thoracic surgeon

## <u>The Lung</u>

#### Embryology


- Bronchial system
- Alveolar system
- Anatomy
  - Lobes
  - Fissures
  - Segments
  - Blood supply

### Airways




© Elsevier Ltd. Drake et al: Gray's Anatomy for Students www.studentconsult.com


# **Bronchopulmonary Segments**



## **Bronchopulmonary Segments**




## **Bronchopulmonary Segments**



# **Blood Supply**

- Lungs do not receive any vascular supply from the pulmonary vessels (pulmonary aa. or veins).
- Blood delivered to lung tissue via the bronchiole arteries.
- Vessels evolve from aortic arch.
- Travel along the bronchial tree.


### **Blood Supply**



© Elsevier Ltd. Drake et al: Gray's Anatomy for Students www.studentconsult.com

# **Blood Supply**

#### **Bronchial Arteries**




# Airways


- Trachea, primary bronchi, secondary bronchi, tertiary bronchi out to 25 generations
- All comprised of hyaline cartilage
- Trachea
  - Begins where larynx ends (about C6)
  - 10 cm long, half in neck, half in mediastinum
  - 20 U-Shaped rings of hyaline cartilage keeps lumen intact but not as brittle as bone
  - Lined with epithelium and cilia which work to keep foreign bodies/irritants away from lungs


# Bronchioles

- First level of airway surrounded by smooth muscle; therefore can change diameter as in brocho-constriction and broncho-dilation
- Terminal
- Respiratory
- 3-8 orders
- alveoli



#### Trachea and Major Bronchi Anterior View






#### Intrapulmonary Airways


©Novartis

## **DISEASES OF THE LUNG**

#### Congenital

- Agenesis
- Hypoplasia
- Cystic adenomatoid malformation
- Pulmonary sequestration
- Lobar emphysema
- Bronchogenic cyst







### A. Lung Abscess

- Causes
- Clinical Features
  - Copious production of foul smelling sputum
- Investigation
  - **C** X R



#### Treatment

- Antibiotics
- Drainage
  - Internal
  - External
- Pulmonary resection
- Indications
  - **1.** Failure of medical RX
  - 2. Giant abscess ( >6cm)
  - 3. Hemorrhage
  - 4. Inability to R. /Out carcinoma
  - **5.** Rupture with resulting empyema
- Type of Resection
  - Lobectomy

B. <u>Bronchiectasis</u> <u>Definition</u> Bronchial dilatation

#### <u>Causes</u>

- Congenital
- Infection
- Obstruction

### **<u>Clinical Features</u>**


- Cough
- Dyspnea
- Haemoptysis (50%)
- Clubbing

#### **Investigation**

- Bronchogram
- CT
- Bronchoscopy

#### **Treatment**


- Medical
  - Resolve most cases
- Surgical
  - Failure of medical Rx
  - Patient with localized disease
  - Cystic type













#### C. <u>Tuberculosis</u>

- \* 30,000 new cases occur annually in U.S.A
- Cause
  - Pulmonary
  - Extra-pulmonary
- Investigation
  C X R





#### Treatment

- Medical
- Surgical
  - Failure of medical Rx
  - Destroyed lobe or lung
  - Pulmonary haemorrhage
  - Persistent open cavity with + ve sputum
  - Persistent broncho pulmonary fistula

### D. <u>Aspergillosis</u>


- Cause
  - Aspergillus fumigatus, A. niger
- Mode of Transmission
- Forms
  - Allergic
  - Saprophytic
  - Invasive
- Saprophytic form
- <u>C-F</u>
  - Aspergilloma
  - Chronic productive cough
  - Haemoptysis (patient with preexisting Disease).


### Investigations

- Skin test
- Sputum
- Biopsy (Invasive)
- CXR

#### Treatment

- Medical
- Surgical
  - Indications
    - A significant aspergilloma
    - Haemoptysis
  - Type of resection
    - Segmentectomy
    - Lobectomy
    - Pneumonectomy





# E. <u>Hydatid cyst</u>


#### <u>Cause</u>


Echinococcus granulosus

<u>Diagnosis</u>

**Treatment** 

















## <u>Tumor</u>

#### ≻<u>Benign</u>

- ≻<u>Malignant</u>
  - Primary
  - Secondary

A. <u>Primary lung carcinoma</u> <u>Incidence</u>

**Risk Factor** 

- Smoking
- Others

**Pathology** 

- 1. Adenocarcinoma
- **2.** Squamous cell carcinoma
- **3.** Large cell carcinoma
- 4. Small cell carcinoma
- NSCLC vs. SCLC

#### **Clinical Features**

- Asymptomatic
- Symptomatic
  - Lung
  - Surrounding structures
  - Rec. L. nerve
  - Oesophagus
  - C8, T1 nerve
  - Sympathetic
  - Pleura
  - SVC

- distal (para-neoplastic syndrome)
  - PTH
  - ADH
  - ACTH
  - Hypertrophic pulmonary osteoathropathy


### **Investigations**

- CXR
- Bronchoscopy
- Trans-thoracic needle aspiration
- CT Scan
- MRI




(see table)


# Tumor




# Tumor








# Bronchoscope



# Tumor



#### lung-cancer-upper-lobe



#### <u>Management</u>

#### **Depends on:**

- Stage
- Cell Type
- Patient Physical fitness


### NSCLC

- Surgical
- Radiotherapy
- Chemotherapy
- SCLC
- Chemotherapy
- Radiotherapy

#### NEW INTERNATIONAL REVISED STAGE GROUPING

Stage 0	TIS
Stage IA	T1, NO, MO
Stage IB	T2, NO, MO
Stage IIA	T1, N1, MO
Stage IIB	T2, N1, MO
	T3, NO, MO
Stage IIIA	T1-3, N2, MO
	T3, N1, MO
Stage IIIB	T4, Any N, MO
	Any T, N3, MO
Stage IV	Any T, Any N, M1

Mountain CF. Chest 1997; 111



# **B. Secondary Lung Carcinoma**

- **Solitary Lung Nodule**
- Primary Carcinoma
- Tuberculous Granuloma
- Mixed tumor
- °2 Carcinoma
- Miscellaneous
- Benign Vs. Malignant


**Hamartoma-Carcinoid** 


- Age
- Sex
- X-ray
  - Size
  - Time
  - Calcification


# THE MEDIASTINUM

#### > <u>Anatomy</u>


- Boundaries
- Divisions
  - Traditional
  - Clinical
- Access: Mediastenoscopy, mediastenotomy
- Mediastinal mass lesions
  - A. Anterior mediatinum(5 T's)
  - **B. Middle Mediastinum(Cyst)**
  - **C.** Posterior mediastinum(Neurogenic)











### **THYMOMA**

### ►<u>Incidence</u>

- The commonest tunmor of A.M.
- Peak 40-60 y.
- M : F (1 : 1)



- Classification
  - Epithelial
  - -Lymphocystic
  - Lymphoepithelial
  - -Spindle cell
- Benign vs. malignant
- Stages
  - I, II, III, IV

### **<u>Clinical Features</u>**


- Asymptomatic
- Symptomatic
  - Mass effect
  - Systemic effect
    - M.G. is the commonest 40-50%

}

#### **Investigation**


- CXR
- CT Scan
- Biopsy
- Bronchoscopy
- Esophagoscopy } cases
- Angiogram

Selected









#### Treatment


- Benign → complete excision
- Malignant → complete excision if possibal

If non-resectable } post-op

- Or } Radiotherapy
- **Resection incomplete** }

### <u>Trauma</u>

- RTA
- Fracture Ribs Simple Complicated
- Haemothorax
- Pneumothorax
- Flail chest
- Lung Contusion and ARDS








# **Flial Chest**

Fracture of several adjacent ribs in two or more places. Flait may be complicated by lung contusion or laceration


Complete sternochondral separation with depression of sternum. Possibility of injury to heart and/or great vessels must also be considered




Accumulation of blood in pleural space

Hemothorax.


















## **Chest Wall**

- Deformity:
  - Pectus excavatum
  - Pectus Carniatum
- Infection
- Chest wall tumor
- Thoracic outlet Syndrome.










### <u>Pleura</u>


- Spontaneous preumothorax
- Pleural effusion
- Empyema
- Mesothelioma .















C5411 W2859









Position of skin incisions, showing camera port and working port anteriorly



Use of a retractor to hold open the working port.

