



Khalid H Al Malki, MD, PhD

Consultant, Associate Professor

Phoniatrics (Voice, Communication and Swallowing Disorders)

Head, Communication and Swallowing Disorders Unit (CSDU)

ENT Department

King Abdulaziz University Hospital

King Saud University, Riyadh, Saudi Arabia.

Head, Communication and Swallowing Division (CSDD)

ORL/HNS Department

Riyadh Military Hospital, Riyadh, Saudi Arabia

<http://faculty.ksu.edu.sa/kmalky/default.aspx>





المملكة العربية السعودية
وزارة التعليم العالي
جامعة الملك سعود
كلية الطب
مستشفى الملك عبدالعزيز الجامعي



**Communication and Swallowing
Disorders Unit (CSDU)**



كرسي بحث
أمراض الصوت والبلع



Internet Site

<http://vas.ksu.edu.sa>



The background features three balloons: a light green one at the top left, a light blue one in the middle left, and a light purple one at the bottom left. Each balloon has a thin streamer and is surrounded by several small, yellow, triangular confetti-like shapes.

Communication and Swallowing Disorders

Khalid H Al Malki, MD, PhD

Three balloons in green, blue, and purple are positioned on the left side of the slide, with yellow triangular rays emanating from them, suggesting a festive or celebratory theme.


Aim of this presentation:

Introduce Communication and Swallowing Disorders and how they are managed generally, and NOT to cover all disorders in details.

Communication Disorders

Communication difficulties have an impact on the following aspects:

- **Academic,**
- **Social,**
- **Psychological,**
- **Employment,**
- **Professional,**
- **Financial,**
- **Family relations.**



Language

A symbolic arbitrary system relating sounds to meaning.

Speech

A neuro-muscular process whereby language is uttered. It includes the coordination of respiration, phonation, articulation, resonance and prosody.

Voice

The result of vibration of the true vocal folds using the expired air.

Swallowing

The process of successful passage of food and drinks from the mouth through pharynx and esophagus into the stomach.

Who is managing Communication and Swallowing Disorders?

Two schools:

- 1. Phoniaticians (MD's).**
- 2. Speech-Language pathologists.**

What is Phoniatics?

- ✱ **A medical specialty that deals with communication and swallowing disorders.**
- ✱ **It stems mainly from ORL (ENT), especially when dealing with VOICE disorders.**

Union of the European Phoniaticians (UEP)

www.phoniatics-uep.org

Phoniatrics (cont.)

☀ In 1931, "Phoniatrics" was officially recognized and acknowledged as an independent medical specialty of its own in Sweden.

☀ In 1992, "Phoniatrics & Pedaudiology" became a new self-standing medical discipline in Germany.

Union of the European Phoniatricians (UEP)
www.phoniatrics-uep.org

Phoniatrics (cont.)

A medical subspecialty of ENT (Otorhinolaryngology) as approved by:

- Saudi Commission For Health Specialties.
- ENT Department, King Saud University.
- Saudi Society of Otorhinolaryngology.

The Phoniatician:

- Deals with the patient from a **MEDICAL** prospective.
- Can perform endoscopies without any limitations.
- Can prescribe medications when needed.
- Can perform surgeries if interested and trained.

JALP

Is Phoniatics recognized internationally?


The International Association of Logopedics and Phoniatics (IALP) is a global organization which promotes the improvement of care for people with communication disorders.

www.ialp.info

IALP:

IALP represents a professional body of:

- ✓ **125.000 members in human communication**
- ✓ **56 affiliated national societies from**
- ✓ **38 countries and**
- ✓ **over three hundred individual members.**
- ✓ **IALP has informative status with UNESCO, UNICEF, WHO,**



Three balloons in green, blue, and purple are positioned on the left side of the slide, with yellow triangular streamers trailing from them.

Language Disorders

Khalid H Al Malki, MD, PhD

I. Language Disorders:

[1] Delayed Language Development (DLD):

Definition of DLD:

Delay or failure to acquire language matched with age.

Three colorful balloons (green, blue, and purple) with yellow streamers are positioned on the left side of the slide.

Prerequisites of normal language development:

- 1. Normal brain function.**
- 2. Intact sensory channels (eg auditory).**
- 3. Normal psyche.**
- 4. Stimulating environment.**

Causes of DLD:

A) Brain damage:

- Diffuse brain damage (MR).
- Brain damaged motorly handicapped child (CP).
- Minimal brain damage (ADHD).

B) Sensory deprivation:

Hearing impairment.

C) Psychiatric disorders:

- Autism.
- Childhood schizophrenia.

D) Non-stimulating environment.

E) Idiopathic.

Diagnosis of the Cause of DLD:

- I. History taking.**
- II. Physical examination.**
- III. Investigations:**
 - Psychometry (IQ).**
 - Audiometry.**

DLD Sheet

Management of DLD:

- ✿ **Early detection.**
- ✿ **Providing the suitable aid (HA or CI).**
- ✿ **Family counseling.**
- ✿ **Language therapy.**

9. Language disorders:

[2] Dysphasia:

Definition:


Language deterioration after its full development due to brain insult: infarction, hemorrhage, atrophy, etc

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Broca's area
Formulates
a speech
response and
stimulates
motor cortex

Motor cortex
Stimulates muscles
that produce
speech

Wernicke's area
Processes
incoming
speech and
comprehends it



Types of dysphasia:

1. Expressive.
2. Receptive.
3. Mixed predominantly expressive.
4. Mixed predominantly receptive.
5. Global.

Diagnosis of dysphasia:

- I. History taking.**
- II. Physical examination: ... , neurological exam.**
- III. Investigations:**
 - CT / MRI brain.**
 - Dysphasia test.**
 - Psychometry (IQ).**
 - Audiometry.**

Dysphasia Sheet

Management of dysphasia:

Individualized:

- **Management of the cause.**
- **Physical rehabilitation (Physiotherapy).**
- **Family counseling.**
- **Language therapy.**
- **Alternative and augmentative communication.**

Three balloons in green, blue, and purple are positioned on the left side of the slide, with yellow triangular streamers trailing from them.

Speech Disorders

Khalid H Al Malki, MD, PhD

Speech disorders:

1. *Dyslalia (Misarticulation):*

Definition:

Faulty articulation of one or more of speech sounds not appropriate for age.

Types of dyslalia:

A) Sigmatism (/s/ defect):-

- Interdental sigmatism.
- Lateral sigmatism.
- Pharyngeal sigmatism.

B) Back-to-front dyslalia:-

/k/ /t/
/g/ /d/

C) Rotacism (/r/ defect).

D) Voiced-to-nonvoiced dyslalia:-

/g/ /k/
/d/ /t/
/z/ /s/ etc...

Diagnosis of dyslalia:

- I. History taking.**
- II. Physical examination: ... , tongue, ...**
- III. Investigations:**
 - Audio recording.**
 - Articulation test.**
 - Psychometry (IQ).**
 - Audiometry.**

Dyslalia Sheet

Management of dyslalia:

- **Treatment of the cause:**
 - . **Tongue tie.**
 - . **Dental anomalies.**

- **Speech therapy.**

II. *Speech disorders:*

2. Stuttering:

Definition:

The intraphonemic disruptions resulting in sound and syllable repetitions, sound prolongations, and blocks.

Normal dysfluency:

- **Less than 6 years.**
- **Only repetitions.**
- **No associated muscular activity.**
- **Not aware.**

Incidence of stuttering: 1%.

Onset:

- **Earliest = 18 months.**
- **Latest = 13 years.**

Epidemiology:

- **more in families with history of stuttering.**
- **can occur in mentally retarded.**
- **very rare in the hearing impaired.**

Gender ratio:

4 : 1 (male : female)

Theories of Stuttering:

The exact cause is unknown.

- Organic theory.**
- Neurosis theory.**
- Learning theory.**

Diagnosis of stuttering:

- I. History taking.**
- II. Physical examination: APA, VPA, ...**
- III. Investigations:**
 - Audio and video recording.**
 - Stuttering severity (eg SSI).**
 - Articulation test.**
 - Psychometry (IQ).**

Stuttering Sheet

Auditory Perceptual Analysis (APA):

A. Core behaviors:

- Intrapophonemic disruption.
- Repetitions.
- Prolongations.
- Blocks.

B. Secondary reactions:

- Muscular activity and struggle.
- Interjection.
- Word substitutions and circumlocution.

C. Concomitant reactions:

- Fear.
- Breathing (antagonism, interruption, prolongation, cessation, ...).
- Eye contact.
- Skin pallor/flushing.

Management of stuttering:

☀ **Family and patient counseling.**

☀ **Speech therapy:**

a. **Indirect therapy: if not aware.**


b. **Direct therapy: if aware.**

Speech disorders:

3. Hypernasality:

Definition:

Faulty contamination of the speech signal by the addition of nasal noise. It results from velopharyngeal insufficiency (VPI).



Causes of hypernasality:

I. Organic:

1. Structural:

a) Congenital:

- Overt cleft palate.
- Submucous cleft palate.
- Non-cleft causes:
 - . Congenital short palate.
 - . Congenital deep pharynx.

b) Acquired:

- Adenotonsillectomy.
- Palatal trauma.
- Tumors of the palate & pharynx.

2. Neurogenic:

- Palatal upper motor neuron lesion.
- Palatal lower motor neuron lesion.

Causes of hypernasality (cont.):

II. Non-organic (Functional):

- Faulty speech habits.
- Mental retardation.
- Neurosis or hysteria.
- Hearing impairment.
- Post-tonsillectomy pain.

Effects of VPI:

- **Feeding problems: nasal regurgitation.**
- **Ear infections (tensor palati: V).**
- **Psychosocial problems.**
- **Communicative problems:**
 - . **Speech: hypernasality.**
 - . **Language: DLD.**
 - . **Voice: hyper or hypofunction.**

Diagnosis of hypernasality:

I. History taking.

II. Physical examination:

- **General.**
- **ENT examination: ... , palate (inspection, palpation) ...**
- **Simple tests:**
 - . **Gutzman's (a/i) test.**
 - . **Czermak's (cold mirror) test.**

III. Investigations:

- **Audio recording.**
- **Fiberoptic nasopharyngolaryngoscopy.**
- **Psychometry (IQ).**
- **Audiometry.**
- **Articulation test.**
- **Nasometry.**

Hypernasality Sheet

Management of hypernasality:

- **Team work.**
- **Feeding.**
- **Hearing.**
- **Maxillofacial.**
- **Palatal and lip surgeries.**
- **Obtulators.**
- **Communication:**
 - . **Language: Language therapy.**
 - . **Speech: Speech therapy.**
 - . **Voice: Voice therapy.**

Speech disorders:

4. Dysarthria:

Definition:

Any combination of disorders of respiration, phonation, articulation, resonance, and prosody, that may result from a neuromuscular disorder.

Types of dysarthria:

1. Flaccid dysarthria:

- **Lesion: lower motor neuron level.**
- **Communication:**
 - * **breathy phonation.**
 - * **hypernasality.**

2. Spastic dysarthria:

- **Lesion: upper motor neuron level.**
- **Communication:**
 - * **strained strangled phonation.**
 - * **labored breathing.**

Types of dysarthria (cont.):

3. Ataxic dysarthria:

- Lesion: cerebellum level.
- Communication:
 - * increased equal stresses.
 - * irregular articulatory breakdown.

Types of dysarthria (cont.):

4. Dyskinetic dysarthria:

- Lesion: basal ganglia level.

A. Hypokinetic type (Parkinsonism):

*** breathy phonation.**

*** rapid rate.**

*** short rushes of speech with final decay.**

B. Hyperkinetic type:

i. Quick hyperkinetic (Chorea):

*** variable rate and loudness.**

ii. Slow hyperkinetic (Athetosis):

*** slow rate.**

Types of dysarthria (cont.):

5. *Mixed dysarthria:*

- may be the most common.

- **Examples:**

- * **Motor neuron disease Flaccid + Spastic.**
- * **Multiple sclerosis Ataxic + Spastic.**
- * **Wilson's disease Ataxic + Spastic + Hypokinetic.**

Diagnosis of dysarthria:

I. History taking.

II. Physical examination: ... , mouth, palate, ... , neurological exam, ...

III. Investigations:

- **Audio recording.**
- **Fiberoptic nasopharyngolaryngoscopy.**
- **CT/MRI brain**
- **Dysphasia test.**
- **Psychometry (IQ).**
- **Articulation test.**
- **Audiometry.**
- **Nasometry.**
- **MDVP.**
- **Aerodynamics (Aerophone II).**

Management of dysarthria:


Individualized:


- **Management of the cause.**
- **Patient counseling.**
- **Communicative therapy:**
 - * **Articulation.**
 - * **Phonation.**
 - * **Resonance.**
 - * **Respiration.**
 - * **Prosody.**
- **Alternative and augmentative communication.**


Three balloons in green, blue, and purple are positioned on the left side of the slide, with yellow triangular streamers trailing from them.


Voice Disorders

Khalid H Al Malki, MD, PhD









Prerequisites of “normal” voice production:

- 1. Normal range of movement of vocal folds.**
- 2. Normal mobility of mucosa on deep layers.**
- 3. Optimal coaptation of vocal folds’ edges.**
- 4. Optimal motor force.**
- 5. Optimal pulmonary support.**
- 6. Optimal timing between vocal fold closure and pulmonary exhalation.**
- 7. Optimal tuning of vocal fold musculature (int. & ext.).**

Three colorful balloons (green, blue, and purple) with yellow streamers are positioned on the left side of the slide.

Definition of dysphonia:

- **“Difficulty in phonation”.**
- **“Change of voice from his /her habitual”.**
- **“Hoarseness” = roughness & harshness of voice.**

Etiological classification of dysphonia:

I. Organic Causes

II. Non-Organic Causes

Habitual

Psychogenic

**III. Minimal Associated
Pathological Lesions
(MAPLs)**

IV. Accompaniment of Neuro-psychiatric Ailments

٩٩٩. Voice disorders:


A) Organic voice disorders:

- . Congenital.**
- . Inflammatory.**
- . Traumatic.**
- . Neurological.**
- . Neoplastic.**
- . Hormonal.**
- . Status post-laryngectomy.**

Sulcus vocalis



Laryngeal carcinoma



Respiration



Phonation

Left vocal fold paralysis



Respiration



Phonation

999.Voice disorders:

B) Non-organic voice disorders:

i. Habitual:

- 1. Hyperfunctional childhood dysphonia.**
- 2. Incomplete mutation.**
- 3. Phonasthenia (Voice fatigue).**
- 4. Hyperfunctional dysphonia.**
- 5. Hypofunctional dysphonia.**
- 6. Ventricular dysphonia.**

Hyperfunctional dysphonia



Respiration



Phonation

Phonasthenia



Respiration



Phonation

Three colorful balloons (green, blue, and purple) with yellow streamers are positioned on the left side of the slide.

B) Non-organic voice disorders (cont.):

ii. Psychogenic:

1- Psychogenic dysphonia.

2- Psychogenic aphonia.

٩٩٩. Voice disorders:

c) Minimal associated pathological lesions (MAPLs) :

- 1. Vocal fold nodules.**
- 2. Vocal fold polyps.**
- 3. Vocal fold cysts.**
- 4. Reinke's edema.**
- 5. Contact granuloma.**

Vocal Fold Nodules: Adult Type



Respiration



Phonation

Vocal Fold Nodules: Juvenile Type




Respiration



Phonation





Left Vocal Fold Polyp with a Reaction



Respiration



Phonation



Respiration



Phonation



Respiration



Phonation


Left Vocal Fold Cyst



Respiration



Phonation




Right-sided Reinke's Edema



Respiration



Phonation



Right-sided Contact Granuloma



Respiration



Phonation



Diagnosis of dysphonia:

- I. History taking.**
- II. Physical examination: APA , ... , neck , ...**
- III. Investigations:**
 - Audio recording.**
 - Digital laryngostroboscopy.**
 - Digital laryngokymography.**
 - Acoustic analysis (MDVP).**
 - Aerodynamic analysis (Aerophone II).**
 - GERD (LPR) work-up.**
 - CT neck.**

***Voice Sheet**

Auditory Perceptual Analysis (APA):

1. **Overall grade:** normal, slight, moderate, severe.
2. **Character (quality):** strained, leaky, breathy, rough.
3. **Pitch:** increased, decreased, diplophonia, normal for age.
4. **Register:**
 - habitual register: modal, falsetto, vocal fry.
 - register break.
5. **Loudness:** loud, soft, fluctuation, normal.
6. **Glottal attack:** hard, soft, normal.
7. **Associated laryngeal functions:** cough, whisper, laughter.




Stroboscopic examination:


- 1- **Glottic closure:** complete, incomplete (shape).
- 2- **Glottic gap:** site, size.
- 3- **Glottic wave:** great, normal, small, absent.
- 4- **Amplitude:** great, normal, small, zero.
- 5- **Symmetry:** in phase, in amplitude.
- 6- **Phase closure:** open phase predominate, close phase predominate.
- 7- **Stroboscopic fixation.**
- 8- **Additional morphological findings.**

VIDEOKYMOGRAPHY (VKG) - Principle


STANDARD mode



HIGH-SPEED (VKG) mode



Time (total c. 18 ms)




Videolaryngokymogram

Videolaryngokymography:

- **Symmetry (both sides) in:** # Amplitude.
Mucosal wave.
Phase.
- **Periodicity (same side) in:** . Amplitude.
. Glottal cycle timing.
- **Closed phase.**

CSL (MDVP)



Aerophone II





C:\AERO\PGG\ELHAMM.AP2


VITAL CAPACITY

FLOW: F1000 5

0.000 SEC

0.000 SEC

20.0



ANTIALIAS
SELECT:
INS + INS
UNSELECT:
DEL
TOGGLE
MARK
ZOOM + Z.
CALCULATE

PGDOWN FORW. PGUP BACKW. +- XSCALE ‡ YSCALE + CURSOR PRINT END



C:\AERO\PGG\ELHAMM.AP2

MAX SUST. PHONATION

R


SPL: 50-100 dB

FLOW: F 300 2.

0.000 SEC

0.000 SEC

20.00



ANTIALIAS
SELECT:
INS + INS
UNSELECT:
DEL
TOGGLE
MARK
ZOOM + Z.
CALCULATE

PGDOWN FORW. PGUP BACKW. +- XSCALE ‡ YSCALE + CURSOR PRINT END

Khalid H Al Malki, MD, PhD

C:\AERO\PGG\SHAWQEEW.AP2

IPIPI (VOICE EFF.)

SPL: 50-100 dB


PRESS: 10 cm H2O

FLOW: F 300 1

0.000 SEC

0.000 SEC

20.0




ANTIALIAS
 SELECT:
 INS + INS
 UNSELECT:
 DEL
 TOGGLE
 MARK
 ZOOM + Z.
 CALCULATE

PGDOWN FORW. PGUP BACKW. +- XSCALE ↓ YSCALE + CURSOR PRINT END

Khalid H Al Malki, MD, PhD

Management of voice disorders:

- **Pharmacological agents.**
- **Surgical procedures (Phonosurgery).**
- **Technical aid devices.**
- **Voice therapy.**




Three balloons in green, light blue, and purple are positioned on the left side of the slide, with yellow triangular streamers trailing from them.


Swallowing Disorders

Khalid H Al Malki, MD, PhD


Phases of normal swallowing:




1. Oral preparatory phase



2. Oral propulsive phase



3. Pharyngeal phase




4. Esophageal phase

Definition of dysphagia:

- **“Difficulty in moving food from the mouth to the stomach”.**
- **“Odynophagia” = painful swallowing due to a disorder of the esophagus.**

Consequences of dysphagia:

- **Dehydration.**
- **Weight loss.**
- **Aspiration pneumonia.**
- **Airway obstruction.**
- **Loss of joy of eating.**



Diagnosis of dysphagia:

I. History taking.

II. Physical examination:

- **General examination.**
- **Language and Speech assessment.**
- **Vocal tract examination.**
- **Neck examination.**
- **Trail feeding.**

III. Investigations:

- **FEES.**
- **VFES (MBS).**
- **GERD (LPR) work-up.**

Dysphagia Sheet




FEES

FEES protocol of evaluation (Langmore, 2003):

- I. Anatomic and physiologic assessment.**
- II. Assessment of food and liquid swallowing.**
- III. Assessment of therapeutic interventions.**


FEES Form



Residue



Penetration



Aspiration



VFES (MBS)

Management of dysphagia:

☀ Oral vs. Nonoral feeding:

Nonoral feeding when:

- a. Aspiration $> 10\%$.
- b. Oral + pharyngeal transit time > 10 sec.

☀ Direct vs. Indirect therapy:

- a. Direct: food or liquid is given to the patient.
- b. Indirect: no food or liquid is given (only saliva).

☀ Compensatory vs. Therapy techniques:

- a. Compensatory: elimination of symptoms but no change in swallowing physiology, such as postural techniques.
- b. Therapy techniques: change of swallowing physiology, such as swallowing maneuvers.

Management of dysphagia:

- **Swallowing therapy:**
 - **Diet modification.**
 - **Postural techniques.**
 - **Swallowing maneuvers.**
 - **Sensory enhancement techniques.**
 - **Motor exercises.**

- **Surgical treatment, eg medialization laryngoplasty.**

- **Medical (Drug) treatment, eg anti-parkinsonism drugs.**

- **Intraoral prosthesis.**

- **Alternative routes of feeding, eg NG tube feeding.**