

Orbit and Oculoplastics

Adel Alsuhaiibani, MD

Goals and objectives

■ Orbit


- Anatomy and evaluation techniques
- Orbital trauma
- Proptosis

■ Lids


- Anatomy and evaluation techniques
- Trauma
- Lesions
- Malpositions

Anatomy


Right orbit: frontal and slightly lateral view




Bones




Orbital Compartments




Sinuses



Blood Supply



Annulus of Zinn



Evaluation

■ 7 P's

- Pain
- Proptosis
- Progression
- Palpation
- Pulsation
- Periorbital changes
- Past medical history

Pain

- Infection
- Inflammation
- Hemorrhage
- Malignant Lacrimal Gland Tumor




Progression Minutes to Hours

- Hemorrhage
- Lymphangioma
- Varix (upon valsalva)



Progression Days to Weeks

- Children: capillary hemangioma, rhabdomyosarcoma, retinoblastoma, neuroblastoma, leukemia
- Inflammatory disease: idiopathic orbital inflammatory disease, thrombophlebitis, thyroid orbitopathy, recurrent inflamed dermoid
- Infection: orbital cellulitis, abscess, cavernous sinus thrombosis
- Trauma, post surgical, hemorrhage: orbital hemorrhage, lymphangioma
- Malignancy: rhabdomyosarcoma, metastatic tumors, granulocytic sarcomas, adenoid cystic carcinoma
- Carotid-cavernous (C-C) fistula



Infection

■ Preseptal Cellulitis

- Vision, motility, pupils, VF, disc are WNL
- globe itself is not proptotic



■ Orbital Cellulitis

- 90% secondary to sinus disease
- high risk of morbidity and mortality
 - orbital abscess
 - brain abscess
 - cavernous sinus thrombosis



Allergic Eyelid Swelling





Progression Months to Years

- Dermoid cysts
- Benign mixed tumors
- Neurogenic tumors
- Cavernous hemangioma
- Lymphoma
- Fibrous histiocytoma
- Osteoma
- Lipoma
- Glioma
- Meningioma

Proptosis

- Primary orbital neoplasms usually unilateral
- Bilateral proptosis seen in inflammatory, immune processes or systemic diseases

Proptosis


- Inflammatory
 - Thyroid disease – most common cause
 - Orbital pseudotumor
 - Wegener granulomatosis
- Infection (orbital abscess, cellulitis)
- Vascular
 - Orbital hemorrhage
 - Lymphangioma (sudden)
 - C-C fistula
 - Orbital varices-proptosis with Valsalva
- Tumor
 - Benign: cavernous hemangioma, lymphangioma
 - Malignant: adenoid cystic carcinoma, lymphoma, glioma
 - Contiguous: sinus, intracranial nasopharynx, skin
 - Metastatic - lymphoma, leukemia, neuroblastoma
 - Rhabdomyosarcoma

Inflammation


■ Graves disease

- Most common cause of unilateral or bilateral proptosis
- May occur with any thyroid status
- Eye disease not controlled by thyroid ablation
- Treatment options
 - steroids
 - radiation
 - optic nerve decompression





Inflammation



- Idiopathic orbital inflammation
 - orbital pseudotumor
 - myositis
 - prompt response to steroids
 - OU or systemic → think vasculitis (*except in kids)
- Sarcoidosis
 - lacrimal gland
- Vasculitis
 - GCA, PAN, SLE, Wegener's granulomatosis

Lymphoproliferative Disorders


- Lymphoid hyperplasia and lymphoma
 - 20% of all orbital mass lesions
 - salmon patch appearance
 - molds to orbital structures
 - 50% arise in lacrimal fossa
 - 17% bilateral
- Plasma cell tumors
- Histiocytic disorders
 - macrophage based d/o




Proptosis


- Axial
- Non- axial
- Pulsital

Proptosis






■ Wilford.mpg





Pseudoproptosis



Palpation



Pulsation

■ Clinical correlation


- With bruits
 - Cavernous carotid fistula
 - Orbital arteriovenous fistula
 - Dural arteriovenous (a-v) fistula
- Without bruits
 - Meningoencephaloceles
 - Neurofibromatosis
 - Orbital roof defect (condition after surgical removal of orbital roof, sphenoid wing dysplasia)



Periorbital Changes



Rhabdomyosarcoma



- Most common primary orbital malignancy of childhood
- Average age: 7-8
- Sudden onset and rapid evolution of unilateral proptosis
- 90% survival




Past Medical History

Imaging options


- Plain films
- CT scan
- MRI
- Ultrasound

Plain films


- Quick
- R/o foreign bodies
- Infrequently used



Caldwell's view



Base view



Waters' view

CT Scan

■ Strengths

- spatial resolution
- bone
 - fractures
 - bone destruction
 - calcification
- quick- emergencies
 - trauma
- cheaper



CT Scan

■ Weakness


- radiation: 1-2 cGy
- soft tissue definition
- contrast iodinated
 - allergy
- may need MRI anyway
 - (not cheaper)

■ Protocols

- axial and coronal
- +/- contrast



Describe the study



MRI

■ Strengths

- Tissue
 - T1→anatomy
 - T2→pathology
- No radiation

■ Weaknesses

- magnetic
 - pacemakers, surgical clips
- claustrophobia




MRI

■ Protocols

- Axial/coronal/sagittal
- Gadolinium contrast
 - non-iodinated
 - allergies RARE
- orbital lesions
 - fat suppression



Name the study




T1 or T2?

Axial/coronal/sagittal ?

Contrast ?

Lesion ?

Orbital Echography




- Dynamic
- Less expensive +/-
- Availability variable


Facial trauma and fractures

- Midfacial fractures
- ZMC fracture
- Wall and floor fractures
 - medial wall- lamina papyracea
 - orbital floor- blow out vs rim involvement
 - lateral wall and orbital roof- less common
- Optic canal fractures
 - traumatic optic neuropathy


LeForte Fractures



Zygoma





A





B



ZMC Fractures




Floor Fractures



Find the fracture




Optic Canal




May be with or without displaced bony fragments




Lacrimal


















Lacrimal Gland Masses

- Inflammatory
 - Sarcoidosis
 - Orbital Pseudotumor
 - Vasculitis
- Non-inflammatory
 - Lymphoproliferative
 - Epithelial neoplasms




Pleomorphic adenoma

Lacrimal gland fossa lesions

Orbital pseudotumor	duration days to chronic	painful- yes	Ultrasound reflectivity: low	CT: localized or diffuse, molds to bone and globe	Management: systemic steroids, XRT
lymphoma	months	no	low	homogenous, oblong, molds to globe/bone	XRT, CTX (systemic disease)
pleomorphic adenoma (benign mixed tumor)	often > 1 year	no	medium to high, regular internal structure	well circumscribed, globular, possible bony expansion or excavartion	complete excision with capsule without biopsy
Adenoid cystic carcinoma, malignant epithelial tumors	< 1 year	yes (perineural invasion)	medium to high, irregular internal structure	round to oval mass with bony erosion	incisional biopsy, await permanent sections; exenteration

Eyelids

- Anatomy
- Trauma
- Lid lesions
- Lid malpositions





Eyelid Trauma

■ Types

- Blunt
- Sharp/penetrating

■ classification

- lid margin
 - not involved
 - involved*

- canthal involved*
- canalicula involved*

*call ophthalmology

Lid Laceration with Canalicular Involvement



Lid margin spared

- Skin and orbicularis only → skin sutures
- FAT protrusion = septum violated
 - DO NOT suture the orbital septum

Blepharitis



Herpes Zoster Ophthalmicus



Lid Lesions

- Sty



- Chalazion




Xanthelasma




Basal Cell

- 90-95% of malignant eyelid tumors
- Lower lid and medial canthal areas
- Nodular and morpheaform types
- Medial canthal lesions can be problematic
- 3% mortality





Nodular




Morpheaform

Squamous Cell

- 40x less common than BCC
- More aggressive
 - perineural invasion
- Most arise from pre-existing lesions
- Variable presentation



Sebaceous adenocarcinoma



- Highly malignant
- 2x more common in upper lid
- Multicentric
- Separate upper and lower lid lesions in 6-8%
- Pagetoid spread

Eyelid Malpositions


- Ectropion
- Entropion
- Blepharoptosis
- Retraction

Ectropion



- Outward turning of lid margin
- Types:
 - Congenital
 - Involutional
 - Paralytic
 - Cicatricial
 - Mechanical



Entropion



- Inversion of the lid margin
- Types:
 - Cicatricial
 - Involutional
 - Congenital
 - Acute-spastic



Trichiasis



Blepharoptosis

- Drooping or inferior displacement of the upper lid
- Classification:
 - Congenital vs acquired
 - Myogenic, aponeurotic, neurogenic, mechanical, or traumatic
- Evaluation

Myogenic ptosis

- Congenital
 - Dysgenesis of levator
- Acquired
 - Localized or diffuse disease
 - Muscular dystrophy
 - CPEO
 - MG
 - Oculopharyngeal dystrophy



Aponeurotic





- Most common form of ptosis
- High lid crease with normal levator function


Neurogenic

- Acquired and congenital forms
- Acquired:
 - 3rd nerve palsy**
 - Horner syndrome
 - Myasthenia gravis



Levator Function







Treatment



- Mild ptosis, good levator function:
Mullerectomy
- Any ptosis, reasonable levator function:
Levator resection
- Severe ptosis, poor levator function:
Frontalis suspension

Mullerectomy












A

Dermatochalasis



B

Brow ptosis



Brow ptosis

Dermatochalasis



Dermatochalasis



PREOP




POSTOP

Abnormal Eyelid Movements

- Blepharospasm
- Hemifacial spasm
- 7th nerve palsy

Blepharospasm

- Involuntary tonic, spasmodic contraction of orbicularis
- dermatochalasis- rubbing
- brow ptosis- frontalis spasm
- blepharoptosis- levator dehiscence
- ectropion/entropion
- dry eye



Hemifacial Spasm

- Intermittent contractions of the entire side of face
- Present during sleep
- Compression of 7th nerve at the level of the brain stem
- MRI evaluation


Hemifacial Spasm


- Intermittent contractions of the entire side of face
- Present during sleep
- Compression of 7th nerve at the level of the brain stem
- MRI evaluation

7th nerve palsy

- Location of lesion:
 - Supranuclear, brain stem, peripheral
- Cause of paralysis:
 - Bell's
 - Infection
 - Infarct
 - Demyelination
 - Neoplasm
 - Trauma
 - Miscellaneous

Course of the 7th Nerve







Botox in Ophthalmology

Botulinum Toxin


- Clostridium botulinum
- Neurotoxin types A,B,C1,D,E,F,G
- Botox = Botulinum Toxin A
- Blocks the release of acetylcholine
- Onset 3 days
- Peak effect 1-2 weeks
- Duration 6-12 weeks




Blepharospasm






Strabismus



Glabellar Botox





Botox for Crow's-Feet



Thank you for your time and
attention