

# THE PHARYNX

# THE PHARYNX

- **Skull base:**

- Cricoid cartilage anteriorly

- Inferior border of C6 posteriorly

- *Widest portion at hyoid ( 5 cm)*

- Narrowest portion at caudal end (1.5 cm)*

- **Divided into 3 parts:**

- *Nasopharynx*

- *Oropharynx → you can see it directly*

- *hypopharynx*

- **Nasopharynx :**

- *Respiratory function*

- *Anterior : choana (posterior nasal aperture)*

- *Posterior : superior constrictor muscle*

- *Superior : basilar portion of occipital bone*

- *Inferior : soft palate*

- **Oropharynx :**

- *Respiratory and digestive function*

- *Anterior : anterior tonsillar pillar*

- *Posterior : superior and middle constrictor*

- *Superior : soft palate*


- *Inferior : base of tongue , superior epiglottis*

- *Laterally:*

- palatoglossal

- palatopharyngeal arches

- parapharyngeal space



- **Hypopharynx :**

- *Digestive function (narrowest area of digestive canal ,This sphincter is called the cricopharyngeus muscle)*
- *Lies posterior to the larynx*
- *Superior :superior border of epiglottis and pharyngoepiglottic folds*
- *Inferior : inferior border of the cricoid*
- *Posterior/ lateral : middle and inferior constrictors , bodies of C4-C6*
- *Anterior : laryngeal inlet*

- **Pharyngeal Wall:**



- 1-Mucous membrane

- *Nasopharynx*  
*Ciliated columnar epithelium*
    - *Oro and hypopharynx*  
*Stratified squamous epithelium*

- 2-Submucosa

- *Nerves, blood vessels, and lymphatics*
    - *Mucous and salivary glands*
    - *Subepithelial lymphoid*

tissue→



Characteristics of *Waldeyer's Ring*:

- *No afferents*
- *Efferent to deep cervical nodes*
- *No capsule except the palatine tonsils*

### *3-Muscular layer*

*External circular and internal longitudinal (opposite in reminder of GI tract)*

•**External** : 3 constrictors, constrict the wall of the pharynx during swallowing

•**Internal**:

- Elevate pharynx and larynx during speech /swallow

1. Palatopharyngeus

2. Salpingopharyngeus

3. Stylopharyngeus

4. Levator veli palatine

**\*\*There is a weak area in the wall of the pharynx covered by mucosa only called Killian's triangle. It is the area for Zenker's diverticulum**

-Tenses soft palate and opens Eustachian tube during yawn/swallow  
Tensor veli palatini (V3)

-Approximates tongue and soft palate Palatoglossus (CN XI via X)

4-Fibrous layer (Buccopharyngeal fascia) → pharyngeal plexus of nerves and veins.

- **Nerve Supply:**

- Motor Muscles of Pharynx & palate : **Acessory nerve** (cranial) via **Vagus nerve**

**Except:**

- tensor palatini muscle by mandibular division of **trigeminal nerve** (CN V)

- Stylopharyngeus muscle by **glossopharyngeal nerve** (CN IX)

- Secretomotor, Sensory & Taste sensation through Vagus & Glossopharyngeal nerves

- **Blood Supply Of Pharynx :**

- *Arterial : All from External Carotid Art*
- Ascending Pharyngeal artery (*Ext Carotid*)
- Tonsillar Artery (*Facial Art*) *imp!*
- Descending Palatine artery (*Maxillary Art*)
- Ascending Palatine artery (*Facial Art*)
- Branch of *Lingual Artery*

- **Venous :** *Pharyngeal & Pterygoid plexus Common Facial Internal Jugular Vein*

- **Pharyngeal Lymphatic Drainage :-**

- *Nasopharynx : retropharyngeal space*
- *Oropharynx : parapharyngeal space*
- *Hypopharynx : neck*
- *Deep cervical (jugular) nodes*

-Side Notes : when a patient comes with Unilatera cervical lymphadenopathy with sudden conductive hearing loss and middle ear effusion.. Think of **Nasopharyngeal carcinoma** unless proven otherwise.  
-How to confirm ? CT, FNA  
-First presentation of nasopharyngeal carcinoma is streaks of blood in oropharynx.

- **Physiology of the Pharynx:-**

- Breathing (inspiration and expiration)*
- Speech (expiration)*
- Deglutition*
- Swallowing (exproation) 2000/day*
- Ventilation (ME)*
- immunity→ Production of immunoglobulins, plasma cells and lymphocytes*

- ***DISEASES OF THE PHARYNX***

**1-DISEASES OF THE NASOPHARYNX:**

- **ACUTE INFECTION OF NASOPHARYNX**

- *Pathologically: is a part of acute rhinitis (common cold)*
- *Clinically: has no specific clinical features*

- **ADENOIDS**

- *Lymphoid tissue in nasopharynx*
- *Enlarged during childhood*
- *Just lateral and posterior to Torus tubarius is a common area for nasopharyngeal carcinoma.*

- **CLINICAL FEATURES**

- *Nasal obstruction*

- *Mouth breathing*

- *Snoring, sleep disturbance, apnea etc*

- *Ear symptoms due to Eustachian tube obstruction*

- *Nasal tone ( because of hyponasality)*

- *Bilateral OME ( because Hypertrophy of adenoid + the tubal lymph nod)*

- *Adenoid face*

- *Mickey Mouse*

- *Overbite*

- *Long face*

- *Crowded incisors*

- **PreOp Evaluation of adenoid disease:**

- *Triad of:*

- *Hyponasality*

- *Snoring*

- *Mouth breathing*

- *Rhinorrhea*

- *Nocturnal cough*

- *Post nasal drip*

- **Treatment : Adenoidectomy**

- *Local Contraindication of Adenoidectomy→*

*-Palatopharyngeal incompetence*

•*Another Contraindication of Adenoidectomy→*

*-Bleeding*

*-Recent infection*

○ **Adenoidectomy**

✓ **Absolute Indications**

*-Adenoid hyperplasia resulting in sleep disturbances or sleep apnea*

*associated with cor pulmonale*

*-Nasal obstruction associated with orofacial abnormalities*

*- Failure to thrive (not attributable to other causes)*

✓ **Relative Indications**

*-Chronic adenoiditis with persistent sore throat, halitosis, or cervical adenitis*

*-Swallowing difficulties (not attributable to other causes)  
recurrent or chronic otitis media or sinusitis*

✓ **Contraindications**

*-Palatal clefting*

*-Velopharyngeal insufficiency (hypernasal speech, nasal regurgitation)*

✓ **Complications**

*-Immediate and Delayed Hemorrhage*

*- Postoperative Airway Compromise*

*- Dehydration typically secondary to pain*

*-Atlantoaxial (C1-C2) Subluxation*

*-Velopharyngeal insufficiency*

*-Nasopharyngeal Stenosis*

*-Eustachian Tube Dysfunction*

## ***2-DISEASES OF THE OROPHARYNX:***

- 1. Acute tonsillitis*
- 2. Acute non-specific pharyngitis*
- 3. Acute diphtheria*
- 4. Infectious mononeuclosis*
- 5. Vincent 's angina*
- 6. Scarlet fever*
- 7. Moniliasis*

### ***1-ACUTE TONSILLITIS:***

- *Tonsils size : ( occupation of oropharynx )*

#### **Grading system:**

- A.      **0 tonsils in fossa**
- B.      **+1 tonsils less than 25%**
- C.      **+2 tonsils less than 50%**
- D.      **+3 tonsils less than 75%**
- E.      **+4 tonsils greater than 75%**

*N.B. Tonsils size is not an indication for surgery unless it is symptomatic*

- **ETIOLOGY**

- *A disease of childhood, with a peak incidence at about 5 to 6 years of age.*

- *%70viruses →Influenza, Rhinovirus, Adenoviruses, Coronaviruses*
  - *%20bacteria →: GABHS,Strept pneumonia, H. infleunzae, Staph. aurius*
  - *%40are beta-lactamse producing bacteria*

- ***GABHS : Group A Beta Hemolytic Streptococcus.** The most common bacteria causing tonsillitis → Very sensitive to penicillin*

- **Clinical features**

- Dysphagia*
  - odynophagia*
  - Otalgia*
  - Headache*
  - Painful cervical lymphadenitis*




- Fever
- Exudates
- Absence of cough coryza and hoarseness
- ± sever sore throat , deep cervical lymph node enlargement

- **THROAT EXAMINATION**

A. Parenchymatous tonsillitis

B Follicular tonsillitis

C. Membranous tonsillitis



- **NECK EXAMINATION**

- Enlargement and tenderness of the jugulo-digastric lymph nodes

- **INVESTIGATIONS**

- Throat swab
- CBC

- **TREATMENT**

- Symptomatic & supportive treatment
- Antibiotics
  - Penicillin V for 5-7days→ first line
  - Erythromycin – second line
- Steroids
- IV antibiotics
- Recurrent , chronic tonsillitis or obstruction → Tonsillectomy

- **COMPLICATIONS OF ACUTE TONSILLITIS**

**General**→ due to toxin produce by GABHS

- *Acute rheumatism*
- *Acute glomerulonephritis*
- *Septicaemia*

**Local:**

- *Peritonsillitis & peritonsillar abscess ( Quinsy)*
- *Neck Abscess*
- *Parapharyngeal abscess*
- *Retropharyngeal abscess*

- **PERITONSILLAR ABSCESS (QUINSY)**

• *An abscess between the tonsil capsule and the adjacent lateral pharyngeal wall*

- **CLINICAL FEATURES**

- *More common in adults*
- *Usually unilateral*
- *Usually follow an attack of tonsillitis*
- *Sever pain > one side*
- *Unilateral earache and cervical lymphadenitis*
- *More odynophagia & drooling*
- *Trismus*
- *Thickened speech (hot potato voice)*

- **TREATMENT**

- *IV antibiotics*
- *Incision and drainage followed by elective tonsillectomy 6 -8 weeks later.*
- *Hot (abscess) tonsillectomy*

- **Th parapharyngeal space (PPS)**

- *Cone shaped at the lateral wall of pharynx.*
  - Base at temporal bone
  - Apex at the hyoid bone
- *Between*
  - Pharyngeal
  - Lateral and medial pterygoid muscles
- *Most frequently involved with infection .*
- **Contents :** *they are very important because any infection in the tonsils or pharynx can spread and cause parapharyngeal abscess so you should worry about these structures . Local spread can happen also.*

*-Loose fibrofatty tissues*

*- Carotid artery*

*-Internal jugular vein*

*-Cranial nerves IX, X, XI , and XII - Cervical sympathetic chain*

*-Lymph nodes*

■ *Nasal cavity , paranasal sinuses*

■ *Nasopharynx and oropharynx*

■ *Mastoid tip*

- **Communication-:**

*-Submandibular*

*-Retropharyngeal*

*-Parotid spaces*

*-Masticator*

*-peritonsillar*

- **Clinical features**

*-Systemic manifestations*

*-Pain, trismus, swelling*

- **INVESTIGATION**

*-Laboratory and bacteriology*


*-CT*

*-MRI*

- **PRINCIPLES OF TREATMENT**
  - *Secure the airway*
  - *Antimicrobial therapy*
  - *Surgical drainage*
    - *External cervical incision*
    - *In order to avoid injury to the great vessels*
- **Retropharyngeal abscess**
  - ***Between***
    - *Prevertebral fascia*
    - *Posterior pharyngeal wall and oesophagus fascia*
    - *From Skull base to tracheal bifurcation*
    - *Major route → mediastinum*
    - ***Contents*** : *Lymph node (<5 y) that receive from:*
      - *Nose-*
      - *Nasopharynx*
      - *Paranasal sinuses*
      - *oropharynx*
      - *Middle ear*
- **ACUTE RETROPHARYNGEAL ABSCESS** → *Due to suppuration of the retropharyngeal lymph nodes present in the retropharyngeal space*
- **CLINICAL FEATURES**
  - *Systemic manifestations*
  - *Respiratory obstruction*
  - *Odynophagia & Dysphagia*
  - *Swelling of posterior pharyngeal wall (usually unilateral)*
- **INVESTIGATION**
  - *Laboratory and bacteriology*
  - *Plain X-rays*
  - *CT*
  - *MRI*

## ○ TREATMENT OF ACUTE RETROPHARYNGEAL ABSCESS

- Secure airway
- Antimicrobial
- Surgical drainage
- Trans oral



## ○ Complications of peritonsillar , para and retropharyngeal infections

- Respiratory obstruction
- Spontaneous rupture (inhalation pneumonia)
- Extension of infection
  - Other spaces
  - Carotid & internal jugular
  - Mediastinitis

## ○ CHRONIC RETROPHARYNGEAL ABSCESS

- Tuberculous (cold abscess)
- Usually due to TB spines but may be secondary to TB lymphadenitis
- Symptoms are insidious
- Treatment is by anti tuberculous medication, repeated aspiration and external drainage

### ○Ludwig's Angina

- Infection of the submandibular space (floor of mouth)
- Usually secondary to dental infection or trauma

### •TREATMENT

- Secure airway
- Most cases respond to antibiotics ( aggressive one)
- Drainage may be needed

## ***2- Acute non-specific pharyngitis***

- None specific infection

- Causes**

  - Viral (very common) , Bacteria & Fungal

  - None infective : GERD, smoking ,any irritants

- Signs and Symptoms :**

  - fever

  - sore throat

  - Odynophagia "painful swallowing" & dysphagia.

  - drilling of saliva

  - Abscess in the area of the muscle cause spasm, Painful opening of the mouth

  - "tresmus", Lock jaw w/o pain

  - snoring.

**Treatment :** treat the underlying cause , symptomatic, antibiotic

## ***3-Acute diphtheria***

- Caused by *Corynebacterium diphtheria*

- Very rare nowadays because of vaccination

- **PATHOLOGY**

  - Local grayish membrane (composed of fibrin, leukocytes, and cellular debris)

  - Powerful Exotoxins travels to heart and nervous system

- **CLINICAL MANIFESTATIONS**

- Systemic symptoms due to the exotoxins

- Toxemia
- Mild fever
- Tachycardia
- Paralysis

- *Local manifestations*

- Sore throat
- Membrane
- Marked lymphadenitis ('bull neck')imp...

- **TREATMENT**

- *Notification , Isolation ,Rest , Antitoxin serum as early as possible*
- *Antibiotics (erythromycin, penicillin G, rifampin, or clindamycin)*

- **COMPLICATIONS**

- Respiratory obstruction
- Heart failure
- Muscular paralysis

#### **4- INFECTIOUS MONONUCLEOSIS:**

- *Systemic infection caused by Epstein-Barr Virus (EBV)*
- *(Epstein — Barr virus is related to nasopharyngeal carcinoma)*
- *Clinical disease is usually seen in young adults*

- **CLINICAL MANIFESTATIONS**

- Fever
- Pharyngitis and/or tonsillitis
- fatigue
- cervical Lymphadenopathy "bull neck" - Ulceration
- Jaundice

-Atypical lymphocytes

-if we give hem/her Ampicillin it will cause RASH

-Other clinical findings

Splenomegaly – 50%

Hepatomegaly – 10%

Rash – 5%

- **Diagnosis :** *Paul-Bunnell test & Monospot test*

- **Treatment**

- "supportive" we can give steroid for severe cases.

-Avoid ampicillin

- **COMPLICATIONS**

-Autoimmune hemolytic anemia

-Cranial nerve palsies

-Encephalitis

-Hepatitis

-Pericarditis

-Airway obstruction

## **5- VINCENT'S ANGINA**

- Subacute infection due to *Spirochaeta denticolata* and Vincent's fusiform bacillus. ( Acute oropharyngeal ulcerative)
- Poor oral hygiene (fetid breath) → malnutrition→fatigue
- pseudomembranous ulceration
- cervical lymphadenopathy
- penicillin ,metronidazole and local oral hygiene

## **6- SCARLET FEVER**(الحمى القرمزية)

- Acute specific —*Strept haemolyticus*
- Erythroxin cause rash
- Treatment: penicillin





## ○ FUNGAL PHARYNGITIS

### • CAUSES


- Long term antibiotics
- Immunosuppression (Leukopenia, Corticosteroid therapy..etc)
- CANDIDIASIS (MONILIASIS, THRUSH)

### • Thrush

- fungal infection candidiasis is the most common.
- Pseudomembranous candidiasis (Thrush)
- DM or immunodeficiency

### • Treatment

- Nystatin
- Fluconazole



## ○ CHRONIC TONSILLAR HYPERTOPHY

### • CAUSES

- Chronic or frequent acute infections
- Idiopathic (exaggerated immune response)

### • PRESENTATION

- Upper airway obstruction
  - Mouth breathing, snoring
  - Disturbed sleep and apnea
- Pulmonary hypertension, cor pulmonale and heart failure.

### • TREATMENT

- Tonsillectomy & adenoidectomy

- **CHRONIC NON-SPECIFIC PHARYNGITIS**

- *Primary*
- *Secondary*
  - Sinonasal disease*
  - Dental infections*
  - Chest infections*
  - Smoking*
  - Gastro esophageal reflux*

- **CLINICAL FEATURES**

- Sore throat*
- Irritation*
- Cough*
- O/E*

- **TREATMENT**

- *Treatment of the cause*
- *Humidification*

- **CHRONIC SPECIFIC PHARYNGITIS**

- *Tuberculosis*
- *Syphilis*
- *Lupus vulgaris*
- *Leprosy*
- *Sarcoidosis*

- **CHRONIC TONSILLITIS**

- *Persistent or recurrent sore throat*
- *Persistent cervical adenitis*
- *Halitosis*
- *Congested tonsils*
- **TREATMENT** → *Tonsillectomy*

## ○ TONSILLECTOMY

### • INDICATIONS

#### ✓ *Absolute Indications*

- *Obstructing tonsillar enlargement* → resulting in sleep disturbances or sleep apnea associated with cor pulmonale >> IMP
- *Suspected malignancy* (unilateral tonsillar hypertrophy) >> IMP!
- *Tonsillitis resulting in febrile convulsions* (may require a Quinsy tonsillectomy)
- *Persistent or recurrent tonsillar hemorrhage*
- *Failure to thrive* (not attributable to other causes)

#### ✓ *Relative Indications*

- *Recurrent acute tonsillitis* (documented 7 infections in 1 year, 5 infections in 2 years)
- *Peritonsillar abscess*
- *Chronic tonsillitis* with persistent sore throat, halitosis or cervical adenitis
- *One attack of quinsy*
- *Swallowing difficulties* (not attributable to other causes)
- *Tonsillolithiasis*
- *Orofacial or dental disorders* (results in a narrow upper airway)
- *Recurrent or chronic otitis media*
- *Individual consideration*
- *PTA unresponsive to nonsurgical management*

### • CONTRAINDICATIONS

- *Bleeding tendency*
- *Recent URTI*
- *Leukemia hemophilias, agranulocytosis, uncontrolled systemic disease*
- *Relative Contraindications: cleft palate, acute infection*

- **COMPLICATIONS**

- *Hemorrhage*
  - Primary
  - Reactionary
  - Secondary
- *Respiratory obstruction*
- *Injury to near-by structures*
- *Pulmonary and distant infections*

- ✓ **Primary Hemorrhage**

- *Bleeding occurring during the surgery*
- **Causes**
  - Bleeding tendency
  - Acute infections
  - Aberrant vessel
  - Bad technique
- **Management**
  - General supportive measures
  - Diathermy, ligature or stitches
  - Packing

- ✓ **Reactionary Hemorrhage**

- *Bleeding occurring within the first 24 hours postoperative period*
- **Causes**
  - Bleeding tendency
  - Slipped ligature
- **Diagnosis**
  - Rising pulse & dropping blood pressure
  - Rattle breathing
  - Blood trickling from the mouth
  - Frequent swallowing
- **Treatment**
  - General supportive measures

- Take the patient back to OR
- Control like reactionary hemorrhage

✓ **Secondary hemorrhage**

- Occur 5-10 days postoperatively
- Due to infection
- Treated by antibiotics
- May need diathermy or packing

○ **Pharyngeal (Zenker's) Pouch**

- A mucosal sac protruding through Killian's dehiscence
- Most probably related to **neuromuscular incoordination** →
  - Failure of relaxation OR
  - Early closure OR
  - Spasm of cricopharyngeus
- **Clinical Features**
  - Dysphagia
  - Regurgitation
  - Aspiration
- **Diagnosis**
  - Clinical examination
  - Barium swallow
  - Endoscopy
- **Treatment**
  - Excision(surgery)

Done by ENT Team 429  
Mohammed Alfaif