



OBSTETRICS AND GYNECOLOGY

(11) Intra Uterine Growth Restriction (IUGR) + Intra Uterine Fetal Death (IUFD)

Leader: Alanoud Alyousef

Sub-leader: Dana ALdubaib

Done by: Dana aldubaib

Revised by: Ahlam Almutairi

Objectives:

Definitions

Etiology

Management

Prevention

Intra Uterine Growth Restriction (IUGR)

Definition: is failure of a fetus to reach its genetic growth potential

- The best guide is to establish fetal size early in gestation
- Estimated fetal weight can be calculate from **ultrasound measurements** (most accurate for fetal weight= US)
- Abdominal circumference** is affected early in growth restriction owing to decreased glycogen storage in the liver

Low Birth weight and Small for gestation age:

LBW is defined by the WHO as birth **weight < 2.5 kg**, so does not correct for gestation. (**imp after delivery**)

SGA (**Small for gestation age**) is used postnatally to describe a fetus or neonate with growth parameters (e.g EFW, AC, birthweight) below a given centile for gestational age. (**determine by a chart**)

Implications of Growth Restriction

Growth restricted fetuses

- * Have a higher risk of **still birth** and mortality
- * Are most at risk of
 - **hypothermia**
 - **hypoglycaemia**
 - **pulmonary haemorrhage**
 - **infection**
 - **encephalopathy**
 - **necrotizing enterocolitis**
- * Incidence of fetal heart rate abnormalities
- * Higher incidence of operative delivery

Aetiology:

There is a wide range of associations:

o Fetal o Maternal o Placental

These in turn may have a genetic or environmental basis

Fetal causes

a) Chromosomal (Most imp)

Abnormal fetal karyotype can be responsible for up to 20% of growth restricted fetus

- o Early pregnancy, triploidy 58%
- o Trisomy 46%
- o Trisomy 21 and Turners – second trimester

The reason is probably because of lack of cell division or cell growth in either the fetus or placenta

US sign: nasal bone flatness, tricuspid regurge and Nuchal translucency: is a collection of fluid under the skin at the back of your baby's neck. It can be measured using ultrasound: between 11 weeks plus two days and 14 weeks plus one day of pregnancy.

b) Structural anomalies / Structural defects

- Central nervous system
 - Cardiovascular system
 - Gastro intestinal system
 - Genito urinary system
 - Muskilo skeletal
- Are associated with an increased risk of IUGR

c) Infection:

O Malaria – major cause of IUGR – its treatment reduces the incidence of IUGR (area where is endemic)

- o Rubella
- o Cytomegalovirus
- o Toxoplasmosis
- o Syphilis

(Any infection can cause IUGR EVEN Viral)

Can affect cell division and have all been implicated

Nutrient Supply

- o Inadequate maternal nutrition can restrict growth in the 3rd trimester
- o examples are the Dutch Famine and the Leningrad sledge
- o Leningrad sledge 700gm Glucose, amino acids and lactate are the major substrates for the fetus
- o Oxygen : Babies born at higher altitude are smaller than those born at sea level (less oxygen supply)

Placental causes

- * Lack of conversion of spiral arteries into utero placental arteries
- * The low-resistance circulation thus created allows high blood flow to the placenta.
- * In normal pregnancies, end diastolic flow (asses by Doppler) is usually present (umbilical arteries by the early second trimester And increases until term
- * Growth restricted fetuses often have absent or reversed end-diastolic (reversed blood from the fetus to mother BAD sign) Absent > the flow only in systolic which result in hypoxia (Normally the flow for the fetus in systolic and diastolic flow) both bad sign.

Maternal causes

a) Smoking

- Active and passive smoking is a major cause of IUGR
- Such babies weigh between 100-300gm less than other babies
- * > 10 cigarettes/ day is significant (MCQ)
- Male fetus more affected than females
- Mechanism is probably via the higher levels of carboxyhaemoglobin in such fetuses.

b) Alcohol:

Moderate to heavy alcohol consumption can reduce fetal weight by up to 500 gm.

c) Drugs

Heroin and methadone use are associated with growth restriction <490 gm and 280 gm respectively.

d) Chronic Diseases (Most common in our country) any CD

- Congenital heart disease – especially if cyanotic
- Chest disease e.g. cystic fibrosis, bronchitis, kyphoscoliosis and asthma in severe cases where there is marked respiratory compromise.
- Chronic renal diseases – especially if there is hypertension, proteinuria
- Diabetes mellitus – if there is renal disease and vascular disease.

Screening:

Clinical Examination

- Palpation
- Symphyseal – fundal height – higher sensitivities than palpation (To see if it's the same as gestation age)

Ultrasound

- Has a better detection rate for IUGR than clinical examination

Only disadvantage is that the work load will be great if all pregnant women were to be subjected to it too often.

Management

The terms symmetric and asymmetric growth restrictions are descriptive (symmetric all the fetus is small , asymmetrical head is good size and the body is small)

- Growth restriction detected at any gestation without associated anomaly is most likely to represent true growth restriction as a result of utero placental dysfunction
- The earlier the gestation the more likely the fetus is to be aneuploidy or infected.

The gestational age should be checked using the last menstrual period and any early scans.

- Diagnosis of IUGR should be made on serial scans – every 2 weeks
- Thorough survey of the fetus for associated anomalies is undertaken
- Liquor volume should be quantified (amniotic fluid index) by US
- Doppler waveforms of the uterine and umbilical artery should be obtained.

From First Aid:

IUGR

DIAGNOSIS:

First step: Diagnose by confirming serial fundal height measurements with ultrasound.

Second step: Ultrasound the fetus for EFW (although as pregnancy advances, ultrasound fetal weight estimates become increasingly unreliable).

TREATMENT

First step: Explore the underlying etiology and correct if possible.

If near due date: Administer steroids (e.g., betamethasone) to accelerate fetal lung maturity.

Then: Perform fetal monitoring with NST, CST, BPP, and umbilical artery Doppler velocimetry.

A nonreassuring status near term may prompt delivery.

Early –Onset Growth Restriction (<32 Weeks)

The principle differential diagnosis is:

- (a) Chromosomal abnormality or some other genetic problem
- (b) Congenital infection
- (c) Utero placental dysfunction

Findings that would make a chromosomal problem more likely include:

- Normal uterine artery doppler findings
- Normal liquor volume
- Presence of a structural abnormality

The commonest infection associated with IUGR is cytomegalovirus (CMV)

- Mother may have complained of flu-like illness
- Fetus has sonographic findings compatible with CMV (e.g. microcephaly and cerebral calcification).

Utero placental dysfunction is a diagnosis of exclusion, Factors supporting this are:

- A history of growth restriction in a previous pregnancy
- Reduced liquor volume
- Abnormal uterine umbilical artery waveforms

Late-Onset Growth Restriction (>32 Weeks)

Most likely cause is utero placental insufficiency, often associated with the development of pre-eclampsia

Fetal monitoring

Monitoring the growth-restricted fetus involves serial fetal measurement

- Abdominal circumference
- Amniotic fluid index
- Cardiotocography
- Doppler ultrasound

*Fetuses with absent end-diastolic flow are hypoxaemic, these changes may appear up to 5 weeks before demise

*Reversed end-diastolic flow is suggestive of preterminal compromise; the fetus may die within 1-2 days if not delivered.

Amniotic Fluid Index

- Reduction in amniotic fluid index (the sum of the four deepest vertical pools in each quadrant) is associated with an increase in perinatal mortality.

- Fetal urine production is significantly lower in the SGA fetus than in the AGA fetus.
- Decreased renal perfusion results in oligohydramnios

Biophysical Profile

- Breathing
- Tone
- Movement
- Amniotic fluid volume
- Cardiotocography

* Requires about 40 mins observation of fetal breathing movements.
(Takes time not for every fetus only suspect IUGR)

-A persistently abnormal biophysical score is associated with absence of end-diastolic flow

Prevention:

- All women should be encouraged to **stop smoking** since it is the commonest risk factor
- Even passive smoking is harmful – husbands should be persuaded to stop.
- **Early aspirin treatment before 17 weeks** (100-150mg) for patients with previous IUGR babies (possible role of placental thrombosis) **LOW DOSE!**

Labour And Delivery

- In the preterm failure to deliver poses the risk of chronic hypoxia while delivery exposes the neonate to the risks of prematurity
- Most fetuses follow a decompensation cascade:
 - Absent end-diastolic flow > decelerative CTG > reversed end diastolic flow > fetal death
- IUGR fetus is more likely to become more hypoxic in labour
- **With AEDF or reversed EDF, delivery should be by caesarean section**

FROM KAPLAN NOTES

Symmetric IUGR

- All ultrasound parameters (HC, BPD, AC, FL) are smaller than expected.
- Etiology is **decreased growth potential**, i.e., aneuploidy, early intrauterine infection, gross anatomic anomaly.
- Workup should include detailed sonogram, karyotype, and screen for fetal infections.
- **Antepartum tests are usually normal.**

Asymmetric IUGR

- Ultrasound parameters show **head sparing**, but **abdomen is small**.
- Etiology is **decreased placental perfusion** due to chronic maternal diseases (hypertension, diabetes, SLE, cardiovascular disease) or abnormal placentation (abruption and infarction).
- Amniotic fluid index is often decreased, especially if uteroplacental insufficiency is severe.
- **Monitoring** is with serial sonograms, non-stress test, amniotic fluid index, biophysical profile, and umbilical artery Dopplers.

HC= head circumference
BPD= biparietal diameter
AC= abdominal circumference
FL= femoral length

Intra uterine fetal death IUFD

- The term IUFD (Intra uterine fetal death) embraces before before the 28th or 24th week of pregnancy (delayed) miscarriage) and those occurring later which result in **macerated stillbirth**.

- **Maceration** is a destructive process which first reveals itself by blistering and peeling of the fetal skin. This appears between 12 and 24 hours after fetal death. The ligaments are softened and the vertebral column is liable to sag. The skull bones overlap each other at the sutures because of the shrinkage of the brain (**Spalding's sign (loss of tone)**) It takes several days for Spalding's sign to appear after intrauterine death, usually a week or more.

*If with maceration fetus is dead more than 12 h
No maceration less than 12 hour

Causes:

1. One of the commonest is **pre-eclampsia**

Hypertensive spasm of the utero placental vessels which results into reduced oxygen supply to the fetus.

2. **Chronic hypertension**

3. **Chronic nephritis**

Fetus dies from placental infarction and hypoxia even before the age of viability

4. **Hyperpyrexia** – a body temperature over 39.40C can kill the fetus directly (**infection**)

5. **Diabetes** in pregnancy

6. **Fetal malformation**

7. **Placental insufficiency**

8. **Idiopathic**

Management:

1) Conservative – await spontaneous labour

2) Induction of labour

*Prostin E2 (Vaginal pessary)

*IV Nalador

*Oxytocin

* **Exclude coagulation disorder**

- Generally hypofibrinogenaemia does not set in until after about **4 weeks after the IUFD** (the main thing before waiting is normal coagulation profile , wait 4 weeks if not spontaneous labour induce it) if abnormal coagulation > deliver . ***Check fibrinogen to detect DIC**

Causes of Fetal Death Based on Trimester “ from First Aid”

T1 (1–13 WEEKS)

1. Chromosomal abnormalities.
2. Environmental factors (eg, medications, smoking, toxins).
3. Maternal anatomic defects (eg, müllerian defects).
4. Endocrine factors (eg, progesterone insufficiency, thyroid dysfunction, diabetes).
5. Unknown.

T2 (14–27 WEEKS)

1. Anticardiolipin antibodies.
2. Antiphospholipid antibodies.
3. Chromosomal abnormalities.
4. Anatomic defects of uterus and cervix.
5. Erythroblastosis.
6. Placental pathological conditions (eg, circumvallate placentation, placenta previa).

T3 (28 WEEKS–TERM) _ Anticardiolipin antibodies.

1. Placental pathological conditions (eg, circumvallate placentation, placenta previa, abruptio placentae).
2. Infections (eg, toxoplasmosis, CMV,

Management from first aid

- D&E may be used if fetal death occurs in T2. D&E has ↓ maternal mortality compared to PGE₂ labor induction, but also has the risk of uterine perforation.
- Labor induction if fetal death occurs in T3. Induction of labor with vaginal misoprostol is safe and effective even in patients with a prior cesarean delivery with a low transverse uterine scar.
- Every attempt should be made to avoid a hysterotomy.
- The patient should be encouraged to seek counseling due to emotional stress caused by diagnosis of fetal death and length of time between diagnosis and delivery.

Monitoring for infections and coagulation profile if no evidence of problems we can wait for 2-3 weeks but if there is any problem deliver. Examine the fetus for malformations take blood from the cord + fetal tissue + placental tissue

Summery

- IUGR: is failure of a fetus to reach its genetic growth potential
most accurate for fetal weight: US
- LBW is defined by the WHO as birth weight < 2.5 kg, so does not correct for gestation. (imp after delivery)
- Abnormal fetal karyotype can be responsible for up to 20% of growth restricted fetus
- Moderate to heavy alcohol consumption can reduce fetal weight by up to 500 gm.
- Fetuses with absent end-diastolic flow are hypoxaemic, these changes may appear up to 5 weeks before demise
- Reversed end-diastolic flow is suggestive of preterminal compromise ; the fetus may die within 1-2 days if not delivered.
- Early aspirin treatment before 17 weeks (100-150mg) for patients with previous IUGR babies (possible role of placental thrombosis)
- Intra uterine fetal death IUFD One of the commonest causes is pre-eclampsia
- Management of IUFD is waiting UP to 4 weeks for spontaneous abortion BUT before waiting we should check coagulation profile .

MCQ's:

Q1. All these tests can be useful in management of intrauterine fetal growth restriction (IUGR) EXCEPT:

- A. Fetal kick chart
- B. Cardiotocography CTG non stress test
- C. Chorionic villous sampling
- D. Biophysical profile
- E. Umbilical cord Doppler waveforms

Q2. The definition of intrauterine growth restriction (IUGR):

- A. Infant with birth weight below 10th percentile for a give gestation age
- B. Infant with birth weight below 25th percentile for a given gestation age
- C. Infant with birth weight of 2.8 kg
- D. Infant with birth weight below 50th percentile for a given gestation age
- E. Infant born at 30 weeks of gestation with a weight of 2.3 kg

Q3. The following are possible complications of IUGR EXCEPT :

- A. Intra uterine death
- B. Severe hypoxia and fetal distress in labor
- C. Meconium aspiration
- D. Hypoglycemia
- E. Post maturity

Q4. The most serious maternal complication of IUFD:

- A. Acute amnionitis.
- B. Acute psychosis.
- C. Pelvic thrombophlebitis.
- D. Hypofibrinogenemia.
- E. Infertility.

Q5. 38-year-old G5P4 woman at 37 weeks' gestation has a history of preeclampsia in previous pregnancies. She presented to the obstetric clinic for her initial screening at 10 weeks' gestation but was subsequently lost to follow-up. At her initial visit the patient had a blood pressure of 140/70 mm Hg and no proteinuria. Since then, the patient reports experiencing headaches, some visual disturbances, and light-headedness but has not sought health care for any of her symptoms. The patient now presents to the emergency department complaining of a headache. Her blood pressure is 162/90 mm Hg. An ultrasound of the uterus is likely to reveal which of the following?

- (A) Intrauterine growth restriction
- (B) Macrosomia
- (C) Placenta previa
- (D) Polyhydramnios
- (E) Rocker-bottom feet

For mistakes or feedback
Obgynteam432@gmail.com

C
A
E
D
A