

King Khalid University Hospital
Department of Obstetrics and Gynecology
Course 482

**PELVIC INFLAMMATORY DISEASE
(PID)**

PID

- SPECTRUM DISEASE INVOLVE CX, UTERUS, TUBES
- MOST OFTEN → ASCENDING SPREAD OF MICROORGANISMS FROM VAGINA & ENDOCERVIX TO ENDOMETRIUM, TUBES, CONTIGUOUS STRUCTURES
- INCIDENCE ACUTE PID 1–2% OF YOUNG SEXUALLY ACTIVE WOMEN EACH YEAR

ETIOLOGY

- NEISSERIA GONORRHOEAE COMMON CAUSE OF PID
- 85% OF INFECTION → SEXUALLY ACTIVE FEMALE OF REPRODUCTIVE AGE
- 15% OF INFECTION OCCUR AFTER PROCEDURES THAT BREAK CERVICAL MUCOUS BARRIER
- BACTERIA CULTURE DIRECT FROM TUBAL FLUID COMMON :
N. GONORRHOEAE, C. TRACHOMATIS, ENDOGENOUS
AEROBIC, ANAEROBIC, GENITAL MYCOPLASMA SPP.

P I D

- **C. TRACHOMATIS**
 - PRODUCE MILD FORM OF SALPINGITIS
 - SLOW GROWTH (48-72 HR)
 - INTRACELLULAR ORGANISM
 - INSIDIOUS ONSET
 - REMAIN IN TUBES FOR MONTHS/YEARS AFTER INITIAL COLONIZATION OF UPPER GENITAL TRACT
 - MORE SEVERE TUBES INVOLVEMENT

PID

- **N. GONORRHOEAE**
 - **GRAM -VE DIPLOCOCCUS**
 - **RAPID GROWTH (20-40 MIN)**
 - **RAPID & INTENSE INFLAMMATORY RESPONSE**
 - **2 MAJOR SEQUELAE**
 - **INFERTILITY & ECTOPIC PREGNANCY, STRONG ASSO. WITH PRIOR CHALAMYDIA INFECTION**

RISK FACTORS

- **STRONG CORRELATION BETWEEN EXPOSURE TO STD**
- **AGE OF 1ST INTERCOURSE**
- **FREQUENCY OF INTERCOURSE**
- **NUMBER OF SEXUAL PARTNERS**
- **MARITAL STATUS ; 33% → NULLIPAROUS**

RISK FACTORS

- **INCREASE RISK**
 - IUD USER (MULTIFILAMENT STRING)
 - SURGICAL PROCEDURE
 - PREVIOUS ACUTE PID
- **REINFECTION → UNTREATED MALE PARTNERS 80%**
- **DECREASE RISK**
 - BARRIER METHOD
 - OC

DIAGNOSIS

- **COMMON CLINICAL MANIFESTATION**
 - **LOWER ABDOMINAL PAIN 90%**
 - **CERVICAL MOTION TENDERNESS**
 - **ADNEXAL TENDERNESS**
 - **FEVER**
 - **CERVICAL DISCHARGE**
 - **LEUKOCYTOSIS**

DIFFERENTIAL DIAGNOSIS

- ACUTE APPENDICITIS
- ENDOMETRIOSIS
- TORSION/RUPTURE ADX MASS
- ECTOPIC PREG
- LOWER GENITAL TRACT INFECTION

P I D

- **75% ASSO. ENDOCERVICAL INFECTION & COEXIST PURULENT VAGINAL D/C**
- **FITZ-HUGH-CURTIS SYNDROME :**
 - **1-10%**
 - **PERIHEPATIC INFLAMMATION & ADHESION**
 - **S/S ; RUQ PAIN, PLEURITIC PAIN, TENDERNESS AT RUQ ON PALPATION OF THE LIVER**
 - **MISTAKEN DX ; ACUTE CHOLECYSTITIS, PNEUMONIA**

FITZ-HUGH-CURTIS



PID DX

- **CBC**
- **ESR**
- **C-REACTIVE PROTEIN**
- **VAGINAL & CERVICAL SWAB**
- **U/S, CT, MRI**
- **CULDOCENTESIS**
- **LAPAROSCOPIC VISUALIZATION**
 - **MOST ACCURATE METHOD FOR CONFIRM PID**
 - **ALL PT. WITH UNCERTAIN DX, NOT RESPONSE TO RX**
- *** -VE GRAM SMEAR NOT R/O PID**

PID





Table 15.3. Clinical Criteria for the Diagnosis of Pelvic Inflammatory Disease

Symptoms

None necessary

Signs

Pelvic organ tenderness
Leukorrhea and/or mucopurulent endocervicitis

Additional criteria to increase the specificity of the diagnosis

Endometrial biopsy showing endometritis
Elevated C-reactive protein or erythrocyte sedimentation rate
Temperature higher than 38°C
Leukocytosis
Positive test for gonorrhea or chlamydia

Elaborate criteria

Ultrasound documenting tuboovarian abscess
Laparoscopy visually confirming salpingitis

SEQUELAE

• INFERTILITY

- ¼ OF PT HAVE ACUTE SALPINGITIS
- OCCUR 20%
- INFERTILITY RATE INCREASE DIRECT WITH NUMBER OF EPISODES OF ACUTE PELVIC INFECTION

SEQUELAE

• ECTOPIC PREGNANCY

- INCREASE 6–10 FOLD
- 50% OCCUR IN FALLOPIAN TUBES (PREVIOUS SALPINGITIS)
- MECHANISM ; INTERFERE OVUM TRANSPORT
ENTRAPMENT OF OVUM

SEQUELAE

- **CHRONIC PELVIC PAIN**

- 4 TIMES HIGHER AFTER ACUTE SALPINGITIS
- CAUSED BY HYDROSALPINX, ADHESION AROUND OVARIES
- SHOULD UNDERGO LAPAROSCOPE → R/O OTHER DISEASE

- **TOA 10%**

- **MORTALITY**

- ACUTE PID 1%
- RUPTURE TOA 5-10%

TREATMENT

- THERAPEUTIC GOAL
 - ELIMINATE ACUTE INFECTION & SYMPTOMS
 - PREVENT LONG-TERM SEQUELAE

MEDICATION

- **EMPIRICAL ABX COVER WIDE RANGE OF BACTERIA**
- **TREATMENT START AS SOON AS CULTURE & DIAGNOSIS IS CONFIRMED/SUSPECTED**
 - **FAILURE RATE, OPD ORAL ATB → 10–20%**
 - **FAILURE RATE, IPD IV ATB → 5–10%**
- **REEVALUATE 48–72 HRS OF INITIAL OPD THERAPY**

CRITERIA FOR HOSPITALIZATION

TABLE 28.3.

Criteria for Hospitalization of Patients With Acute Pelvic Inflammatory Disease

The following criteria for hospitalization are based on observational data and theoretical concerns:

- Surgical emergencies such as appendicitis cannot be excluded.
- The patient is pregnant.
- The patient does not respond clinically to oral antimicrobial therapy.
- The patient is unable to follow or tolerate an outpatient oral regimen.
- The patient has severe illness, nausea and vomiting, or high fever.
- The patient has a tuboovarian abscess

CDC RECOMMENDED TREATMENT REGIMENS FOR OPD OF ACUTE PID

Table 15.4. CDC Guidelines for Treatment of Pelvic Inflammatory Disease

Outpatient Treatment

Regimen A

Ofloxacin, 400 mg orally 2 times daily for 14 days, or
Levofloxacin, 500 mg orally once daily for 14 days

With or Without:

Metronidazole, 500 mg orally 2 times daily for 14 days

Regimen B

Cefoxitin, 2 g intramuscularly, plus *probenecid*, 1 g orally concurrently, or
Ceftriaxone, 250 mg intramuscularly, or
Equivalent cephalosporin

Plus:

Doxycycline, 100 mg orally 2 times daily for 14 days

With or Without:

Metronidazole, 500 mg orally twice a day for 14 days

CDC RECOMMENDED TREATMENT REGIMENS FOR IPD OF ACUTE PID

Inpatient Treatment

Regimen A

Cefoxitin, 2 g intravenously every 6 hours, or
Cefotetan, 2 g intravenously every 12 hours,

Plus:

Doxycycline, 100 mg orally or intravenously every 12 hours

Regimen B

Clindamycin, 900 mg intravenously every 8 hours

Plus:

Gentamicin, loading dose intravenously or intramuscularly (2 mg/kg of body weight) followed by a maintenance dose (1.5 mg/kg) every 8 hours

TREATMENT

- RX MALE PARTNERS & EDUCATION FOR PREVENTION REINFECTION

- RX MALE PARTNERS → REGIMENS FOR UNCOMPLICATED GONORRHOEAE & CHLAMYDIAL INFECTION
 - CEFTRIAZONE 125 MG IM FOLLOW BY
 - DOXYCYCLINE (100) 1X2^o PC X7DAYS OR
 - AZITHROMYCIN 1GM^o OR
 - OFLOXACIN (300) 1X2^o PC X7DAYS

SURGICAL TREATMENT

- **LAPAROTOMY FOR**
 - SURGICAL EMERGENCIES
 - DEFINITE RX OF FAILURE MEDICAL TREATMENT
- **LAPAROSCOPY**
 - CONSIDER IN ALL PT WITH DDX OF PID & WITHOUT CONTRAINDICATION
 - R/O SURGICAL EMERGENCY
- EVIDENCE OF CURRENT / PREVIOUS ABSCESS
- ACUTE EXACERBATION OF PID WITH BILATERAL TOA

RUPTURED PELVIC ABSCESS

- MORTALITY RATE 10%
- CAN RUPTURE SPONTANEOUS INTO
 - RECTUM
 - SIGMOID COLON
 - BLADDER
 - PERITONEAL CAVITY
- ALMOST NEVER IN VAGINA

THE END

dvdangor@gmail.com