



Ear I-II

1st lecture:

- Gross applied anatomy of the ear
- Nerve supply of the external and middle ears and the principles of referred earache
- Central connection of the vestibulo- cochlear nerve
- Physiology of the external, middle and inner ears

2nd lecture:

- Recognize the congenital anomalies of the external ear
- Diagnose and treat wax accumulation
- Diagnose and treat the common external ear inflammatory conditions
- Discuss the pathology, clinical features and management of AOM

Resources: 436 slides and notes, 435 Team

Done by: Khalid Alhusinan, Mosaed Alnowaiser

Lama Alfawzan, Ghada Alhadlaq, Haifa Alwael

Edited by: Khalid Alhusainan

Safa Al-Osaimi


Revised by: Abdulaziz ALMohammed

All The lectures are based on group A teamwork

Anatomy of the ear

it has 3 parts: Those are simple vedios can summrize the anatomy of [external](#), [middle](#), [inner ear](#) (recommended)

- External ear:** From the outer part till the eardrum (tympanic membrane). It contains the Squamous part of tympanic membrane.
- Middle ear:** (tympanic cavity); From the eardrum till the stapes footplate. It contains the mucosal part of tympanic membrane.
- Internal ear:** Cochlea and vestibule (semicircular canals for angular acceleration and the saccule for linear acceleration). balance. from the vestibule and the three semicircular canals. what is the difference between semi-circular canal and vestibules (contains utricle and saccule) ? semicircular canal is when you are in the elevator "utricle" and you know you are going up and down , also in the car you know it's moving forward and backward " saccule" . for the vestibular its for العباب:)





External ear:

● Formed of Auricles (pinna) and External auditory meatus (auditory canal) and both are lined by skin (Auricle and meatus).

● **Auricle:** is fibrous cartilage "thin" (except lobule area-no cartilage) lined by skin.


● what is its significance?

- In case of Perichondritis (lobule is intact) but in case of any skin problem like Erysipelas, all of auricle is affected. - Auricle is attached to temporomandibular joint (so, movement of this joint will aggravate the pain in case of inflammation of pinna) perichondrium is important. once we separate any perichondrium from a cartilage the cartilage will die (necrosis) so if we have hematoma of the ear (trauma, piercing, etc) and we did not treat it immediately we will end up with necrosis, the septum will separate from the cartilage (the main blood supply to the cartilage) and necrosis of the cartilage will happen.



Perichondritis

(Perichondritis: redness, discharge)



Erysipela

(Erysipelas: skin infection with staph and there is redness)

● **The external auditory meatus (External Auditory Canal (EAC):** (2.5 cm long) is an S shape canal

- it is not straight , if it was water will come in the ear while we are in shower one way to test balance by applying warm air and cold air to the ear if the patient develops dizziness that means that the vestibular function is normal and the vestibular nerve is normal
- some patient will experience dizziness due to changes in temp, vago-vagal attack & cough because some nerves are there like the vagus and glossopharyngeal
- if you didn't pull upward & backward and make it straight you will hit the canal exactly in the (isthmus) which is the narrowest part, where the area of cartilaginous part meets the bony part
- In pediatric it might be straight but in adult it's curved (it's also could be straight if canaloplasty was done)



- (to protect the eardrum and middle ear. Normally we must hump one anterior and one posterior. So, at examination you should pull the auricle posteriorly and superiorly to straighten the canal "Push the pinna upward, backward and lateral". In infant downward and backward.



● **External Auditory Canal consists of:**

○ **lateral third** (outer 1/3) of canal length:


- Cartilagenous
- Contains small amount of subcutaneous tissue
- appendages formed by elastic cartilage and contains ceruminous glands (secrete wax), hair follicles, sebaceous and apocrine glands all together called (apopilosebaceous unit).

* Best places to take cartilage as grafting from, without affecting the shape, are: **TARGUS** (especially in rhinoplasty because it's straight) **Concha and Scaphia** (for tympanoplasty).

* can I take from other places ? yes , for revision surgery rhinoplasty and there is no remaining septal cartilage to use, will take from the ear.

* there are other option that the ear like the costal rip. But ear is better, here morbidity is high


EXTRA PIC



○ **Medial two thirds** (inner 2/3) is osseous:


- Bony
- The narrowest portion is at the bony-cartilaginous junction. No subcutaneous tissue or appendages developed after birth. No hair or wax here! Unless the patient pushes it inside and if so, it won't go out, he must come for wash.

- The skin is thin(0.2mm) skin over bone and easy to be injured during examination. Natural constriction. Another area of constriction is at the tympanic membrane.



● **Anatomical relations of external auditory canal: IMPORTANT**

كثير يحون يشكون من اذانهم والمشكلة تكون ب -
عشان كذا يعتبر واحد (temporomandibular joint) TMJ
لازم الاذن DDX من ال
الباروتيد بتجى انتيروري والفيوريلى-



This is from the lecture only to refresh your memory: Facial nerves gives five branches :temporal and cervical , buccal , zygomatic, (extracranial branches) it comes along the parotid between the superficial and deep lobe from sternomastoid foramen. feel the mastoid . tip on your face there is a groove behind it there were the facial nerves leaves, so a lot of time when we want to do postauricular incision, we will not exceed the tip, if we did we will cause injury to the facial nerve. especially in pediatric (less than 1-2 , who can't move their head yet) the mastoid tip is not long enough, so I must leave 1cm before

the tip

Tympanic membrane (TM):

- it separates the external ear from middle ear
- the Tympanic Membrane is divided into **2 parts**:
 - Pars Tensa. 80%
 - Pars Flaccida. 20% (thin and weak. it goes with negative pressure inside & if you do Valsalva it will go outside)




Figure 1.2 The normal tympanic membrane (left). The shape of the incus is visible through the drum at 2 o'clock. The 'pars flaccida' is the part of the eardrum that covers the upper section of the middle ear. The drum is more 'tense' in the lower section – hence it is called the 'pars tensa'.

- **And consist of three layers:**

- Outer layer stratified squamous epithelium (skin), ectodermal origin. epithelial layer
- The middle layer or lamina propria fibrous layer, mesodermal origin. (present only in pars tensa which makes pars flaccida more prone for perforation) fibrous layer
- The inner layer of endodermal origin, comprising the middle ear mucosa. Mucosal layer

➤ **TM supplied mainly** by V3 (Mandibular) anterior, and X (Vagus) posterior on lateral (outer) aspect, IX (Glossopharyngeal) on medial (inner) aspect.

➤ **How can we determine this is right or left ear?**

By the angle of cone of light and handle of malleus If right-> right ear Left -> left ear.


- In case there was retraction of the tympanic membrane > narrowing of the light cone, and if it was bulging (fusion) > widening of the light cone

Annulus is a fibrous band around the pars tensa that holds the tympanic membrane. if we do tympanoplasty, we do refreshment of the edge then I will take graft and put it under it .why? for epithelization. The graft will prevent the epithelization from going inside making it in one line. So if I did not elevate the annula and fix it again in its place it will be lateralized /blunt and wont give me the normal picture.

Why it's important to be at the same place? because the sound when it stuck to the tympanic membrane and there is vibration it will not transmit to the ossicles. but if the tympanic membrane is thick we will have conductive hearing loss.

if it's affected through marginal perforation that means the (stratified squamous) skin that is inside the external ear canal can go inside and induce a cholesteatoma

Cholesteatoma is not a tumor or high cholesterol, basically it's a normal skin in abnormal place (mucosa.) It will eat the bone. Pars flaccida has no annulus so cholesteatoma can happen through it also.



Middle ear:

- **Lining of the middle ear:**


Mucous membrane: ciliated columnar anteriorly, and cuboidal or flat elsewhere.

Mucous membrane of the middle ear space consists of stratified cuboidal epithelium, which changes to pseudostratified ciliated epithelium around the mouth of the Eustachian tube.

- **Middle ear cleft formed of: important**

- Eustachian (Pharyngo-tympanic) Tube.
- Tympanum (Middle Ear Cavity/proper).
- Mastoid Antrum and Air Cells.

In OR you see opening from middle ear to mastoid is called Aditus (bridge). But From mastoid to middle ear is called antrum (largest air cell in mastoid)



● Eustachian (Pharyngo-tympanic) Tube¹ :

- ◆ Connect the middle ear cavity with nasopharynx “nasal cavity” (upper aerodigestive tract).
- ◆ Lies adjacent to the ICA (internal carotid artery).
- ◆ Normally always closed. But in case of: Yawning, eating, Swallowing (When you swallow sometimes your ear make sound this is ET) → the ET open up by salpingopharyngeus muscle & Tensor tympani
- ◆ when you are at the plane and have URTI, and the pilot landed very fast .so what can we do ? chew, steroid or atropine spary (decongesta when you are at the plane and have URTI,and the pilot landed very fast .so what can we do ? chew, steroid or atropine spery (decongestant nt)
- ◆ open at torus tubarius².
- ◆ **Parts of Eustachian Tube:**
 - Proximal 1/3 is bone.
 - distal 2/3 is fibrocartilaginous, **That is collapsed at rest**
 - Junction between 2 parts is isthmus, **narrowest part of the tube.**

◆ Physiology of Eustachian tube:

- Opens actively by contraction of tensor veli palatine and passively by contraction of levator veli palatine (it releases the tension in tubal cartilage).
- Closed by elastic recoil of elastin hinge + deforming force of Ostmann’s fat pad.
- Protection, Drainage, **Ventilation** “air entry” (**most important function**): The tube protect from anything comes from the nasopharynx to go to the middle ear
- The tube permits aeration of the middle ear and if it is obstructed fluid may accumulate in the middle ear causing deafness.
- The tube equalizes the air pressure during breathing with the external environment
- علشان كذا لما تهبط الطائرة يكون الضغط عالي فنحاول نقله عن طريق العمليات الي تفتح الانبوب هذا منها (Valsalva)
- **The tube is shorter, wider and more horizontal in the infant than in the adult.**
- Secretions or food may enter the tympanic cavity more easily when the baby is supine particularly during feeding.
- The tube is normally closed and opens on swallowing because of movement of the muscles of the palate. This movement is impaired in cleft palate children who often develop accumulation of middle-ear fluid (otitis media with effusion).






Figure 1.3 Diagram to show the anatomy of the middle ear and mastoid air cells



EXTRA 10

Adult vs INFANT

	ADULT	INFANT
Length	36 mm	18 mm
Angle with horizontal	45°	10°
Lumen	Narrower	Wider
Angulation at isthmus	Present	Absent
Cartilage	Rigid	Flaccid
Elastic recoil	Effective	Ineffective
Ostmann's fat	More	Less




ADAM

¹ The Eustachian tube connects the middle ear with the nasopharynx at the back of the nasal cavity. Any fluid in middle ear it will drain into nasopharynx

² **Torus tubarius** of the auditory canal is a mucosal elevation in the lateral aspect of the nasopharynx


❖ Tympanic cavity (Middle ear cavity):

- **Contents of tympanic cavity:** (roof, floor, anterior wall, posterior wall, lateral wall, medial wall.)
 - Ossicles: the malleus, incus and stapes (The smallest bones in the body)
 - Intratympanic muscles: Tensor tympani³, Stapedius⁴
 - nerves: Chorda tympanim, Tympanic plexus.
- stapedius is stronger but they will help each other, if we don't have those two muscle well will all have noise trauma. once you have very strong voice / loudness or frequency higher than 60-70dp, those muscles will contract to prevent the extra-sound from going inside the ear and harm it




في الأعراس لما تخلص نسمع صوت طنين هنا بدت تتأثر التينيسور تحاول تمسك بس للأسف ارتخت في النهاية

- The neck of Stapes receives the insertion of stapedius muscle. Contraction of the stapedius muscle restrict the movement of the stapes (this is considered as a physiologic reflex that protects the inner ear from very loud sounds (Attenuation reflex). - Neck of Malleus receives the insertion of Tensor tympani muscle.
- **Epitympanum:** area above the tympanic membrane is the place where the most acquired cholesteatoma happens because the pars flaccida is here, so when retraction happens this is the first place to get affected called: **Prussak's Space.**
- **Mesotympanum:** area adjust to tympanic membrane, the one we see it once we open tympanic membrane.
- **Hypotympanum:** area below the tympanic membrane



● Nerve supply:

- Sensory nerve supply of the **middle ear mucosa:**
 - Tympanic branch of the glossopharyngeal nerve.
 - Auriculotemporal branch of the trigeminal nerve.
- Motor nerve supply of the **middle ear muscles:**
 - **Stapedius muscle** supplied by the stapedial branch of the **facial nerve.**
 - **Tensor tympani muscle** supplied by the mandibular division of the **trigeminal nerve.**



➤ Clinical importance of walls of middle ear:

- Fracture of temporal bone (roof of middle ear cavity) will be presented by either CSF otorrhea or rhinorrhea.
- Lateral sinus thrombosis secondary to otitis media (posterior wall).
- The middle cranial fossa of the brain is separated from the middle ear by the tegmen tympani.
- 1 st turn of the cochlea forms the promontory
- Chordae tympani is a branch of CN7
- The canal of the carotid a. doesn't go into the middle ear but it's adjacent to it.


³ Attached to neck of malleus has a lowering effect

⁴ Main power for stabilizing sound.

- How many nerves passes through? **Facial, Jacobson (branch of 9th CN), chordae tympani**. Facial pass on top of the stapes, Jacobson passing through promontory, chorda tympani in the middle ear and supply the inguinal nerve for anterior 2/3 of the tongue.
- Facial nerve come from nucleus in pons go **to internal auditory canal along with 8th CN** , then passes into three canals (Labyrinth, tympanic “the most dehiscant [without bone coverage] part of the facial nerve” , mastoid) then it leaves the canal through stylomastoid foramen and turns into 5 branches (temporal, zygomatic, buccal, marginal mandibular, cervical).
- Why this is important? During any ear surgery especially in the stapes, the adhesive part of the facial nerve could be collapsed preventing the surgery⁵


● Anatomical relations of middle ear: **Important!**

- Floor: internal jugular vein and common carotid
- Lateral: tympanic membrane
- Medial: promontory of the cochlea



❖ **Mastoid⁶ antrum and air cells:**

- Air-containing cells of the mastoid process are continuous with the air in the middle ear.
- Pneumatization is complete between the sixth and twelfth years of life.
- Normal tubal function is a prerequisite for biologically active, healthy middle ear mucosa, and thus for the normal process of pneumatization.



- مدخل من الماستويد الي يدخل فيه للمدل اير الفتحة حقتة اسمها antrum

- طيب أنا في المدل اير ويدخل على الماستويد الفتحة الي يدخل فيها اسمها aditus. لما نفحص الاذن ونشوف para flasidis نعيها هذي aditus


- why do we have air cells / sinuses ? to make the skull light ?how sound is transmitted? we have two windows in cochlea oval window & round window. why are they important? the sound wave goes to the tympanic membrane , it moves the membrane and vibrates it (ossicles) , and changes from sound wave to mechanical .then It reach pestim (in the oval window) so it moves it in this way. inside the cochlea there is fluid that is divided into three rooms, upper room is “scala vestibule”, in the middle “scala media”, lower “scala tympani” (it has the round window). so when moves the scala vestibuli the fluid goes into the cochlea swims and goes again to the scala tympani producing some movement in the round window.
- لو ما كانت فيه هذه الازاحة (ازاحة الصوت) كان الفلود ما تحرك من جوا. يعني لو ماكان في مساحة شيء ميكانيكي بحيث أنه يدخل ويطلع. عشان كذا عندنا ويندو to release مساحة الفلود الداخلية كان ما تحرك الي في الوسط الي هو basilar membrane
- what is the basilar membrane? it is a tectorial membrane that has hair cells under it .when the tectorial membrane moves it will help the hair cells like a piano and sound will be produced. when it move I should see a

⁵ المفروض إن الفيشل نيرف يكون مغطى بيوني كنال من المبييتس لين يطلع من برا ، فقط فيه جزء يكون مو مغطى في ٤٠٪ من الحالات ، المشكلة لو كان بورلابسد هنا ما تقدر نسوي العملية

⁶ Mastoid is an air cavity behind the middle ear to mastoid tip, in children less than 2 years it's filled with bone marrow so if we drill there it's going to be bloody. Also, it gets elongated when the child begins to rise his head by sternocleidomastoid muscle.

reflection. This is how we know in surgeries if I fixed the ossicles well or not? we move the ossicles and see a reflection in the round window (like the one we see in the tympanic membrane

➤ **Relationships of the mastoid antrum:**





❖ **SUPPLY OF MIDDLE AND EXTERNAL EAR:**

- corda tympani: is a branch of the facial nerve that gives special sensory for taste anterior 2/3 of the tongue it gives fibers with the lingual nerve and give taste.
- facial nerve : from brain stem (intracranial part) then to the maditus “ internal auditory canal” and then to the temporal bone 3 parts , lastly it exit from stylomastoid and give 5 branches (extra-cranial) .
- **#Those are common questions in the exam#:**
 - what are the 3 parts inside the temporal bone?
 - 1-labyrinthine, in bells palsy usually edema is in the narrowest part witch is here
 - 2-tympanic horizontal segment “ in the middle ear” .
 - 3-mastoid vertical segment “ in the mastoid bone ” .
 - the most common dehiscence of facial nerve? tympanic segment 40% injury to the mastoid is iatrogenic

❖ **sensory supply of middle and external ear:**

- Cervical II & III: **Great auricular nerve** c2, c3 (lobule, lateral/inferior auricle) & **Lesser occipital** c2 (medial surface of pinna)
- V cranial nerve: **Auriculotemporal nerve (V3):** tragus. anterior helix, Ant canal wall
- IX cranial nerve: **Tympanic or Jacobson’s.**
- X cranial nerve: **Auricular or Arnold’s branch of vagus** (concha, Post canal wall) **important** if you put cotton inside the ear you will feel tingling in pharynx this is vagus
- V cranial nerve: **Facial nerve** concha, Post canal wall (if you have infection in facial nerve palsy you have to look in the ear you may see vesicles)



❖ **Referred Earache⁷: important** 50% of ear pain is from outside the ear.

➤ Pain in the ear due to a disease in an area supplied by a nerve that also supply the ear.

- **Cervical II & III:** Cervical spondylosis, neck injury (disc, muscle spasm) etc.
- **V (Trigeminal) cranial nerve:** Dental infections, sinonasal diseases etc.
- **IX (Glossopharyngeal) cranial nerve** (branch of CN 9 called jacobson in the promontory): Tonsillitis, pharyngitis, laryngitis, laryngeal cancer, esophageal foreign body post-tonsillectomy, carcinoma etc.
- **X (vagus) cranial nerve:** Tumors of hypopharynx, larynx & esophagus. One of the signs of recurrence tumors in larynx & pharynx is ear pain. Auriculotemporal nerve (V3): any patient that has dental issue or TMJ, tonsillitis, URTI so when they present with ear pain I have to examine those, dental, pharynx, oropharynx, cervical

Inner ear

● **Consists of:** Osseous Labyrinth and Internal auditory canal.

A) Labyrinth consists of:

○ **Bony Labyrinth** its parts:

- Bony Cochlea 35 mm long, 2.5 turns.
- Vestibule
- Bony semicircular canals.

Its contents:




- **Perilymph fluid:** extracellular-like fluid; found in scala tympani and vestibuli. ($K^{+} = 4 \text{ mEq/L}$, $Na^{+} = 139 \text{ mEq/L}$)

- **Membranous labyrinth:**

- Cochlear duct
- Sacculle (inferior) and utricle (superior) > both form the endolymphatic duct extended to the dura laterally (its important in meniere's disease "increased perilymph" we used it for shunt placement).
- Membranous semicircular ducts.

Its contents:

- **Endolymph:** intracellular-like fluid; found in scala media; contributes to positive DC resting potential of 80 mV in scala media; produced from perilymph by marginal Membranous Labyrinth cells of stria vascularis; absorbed within the endolymphatic sac. ($K^{+} = 144 \text{ mEq/L}$, $Na^{+} = 13 \text{ mEq/L}$)



⁷ During examination of the ear the pt may cough (glossopharyngeal reflex) or vasovagal attack (vagus nerve)

◆ Sensory epithelium:

- Cochlea: **Organ of Corti**⁸ : rests on basilar membrane and osseous spiral lamina; major components include:
 - Outer and inner hair cells.
 - Supporting cells: provide structural and metabolic support.
 - Tectorial membrane.
 - Reticular lamina.
- Utricle & saccule: **maculae**.⁹
- Semicircular canals: **cristae** (angular acceleration). Fluid can move both way that's why responsible for angular acceleration.

◆ Inner canal:

- Consist of cochlea has two window (round&oval) , and semicircular canal
- Membranous part Why its important ? . here is Scala vestibule ,media and tympani
- Inside the membranous part is fluid (endolymph) inside it there is crystal "calcium tubercles " inside each semi-circular canal . They give the sense of going up & down depending on gravity
- Like in BPV(benign positional vertigo) in people who had road traffic accident or prolonged surgery and during transportation the head has been shaken forcefully. Trauma to the head concussion of the inner ear
- فيها مادتين وحدة سايلة وتحتها مادة جيلاينه. البار تكلز موجودة في المادة الجيلاتينية حيث لما تلفين راسك تتحرك بشويش اذا طلعت من المادة الجيلاتينية وصارت تسبح في المادة السائلة. وش بصير؟ مع اي حركة تحسبن ان الدنيا تقلب وتدور مثلا لما تتروش وترفع راسها فوق تقول الدنيا تقلب على ورا أو ولما تنزل تربط الجزمة الدنيا تلف تقعد لها دقيقة كأنها في ملاهي.
- so we do maneuvers to put them back in place (epley maneuvers)
- المانوفرز منتو مطالبين تعرفون كيف تتسوى بس حلو تفهمون معناها, نثبت الراس على حسب السمي سيركلر كنال ونلف 45 درجة وننزلة تحت السرير نشوف اذا عنده سنناقمس او لا العين والاذن مشتبكين مع بعض vestibulo-ocular reflex is very high فلما يصير عندنا اي امبالس للكرستلز نشوف السنناقمس بعده اتهدا ونغير البوزشن نرجعها مكانها لما نعيد التست وماطلعت السنناقمس معناها انها تعدلت
- So its not medical treatment its re-positioning
- Scala media has tectorial membrane and hair cells so with the fluid movement it will move and produce sound
- Each part of the inner ear has sensory organ "important"
- In the utricle and saccule we have organ that moves and cause inflation and deflation with up & down movement its called Macula
- semicircular canal has bulla at the end that had crystal
- the lateral Semicircular canal is the most prominent one so it's the first can be effected by diseases because the most close to external so any ear diseases that eat the bone or cholesteatoma it is the first to be effected.
- So what will happen (for the vestibule) once we have stimulation its goes to the brain stem then to the brain it gives two part one to the spine and one to the oracular muscle
- So we have nystagmus and imbalance this is for the vestibule
- So how can I know that its only vestibular diseases?
 - We know balance is dependent on three thing : proprioception (joint & muscle) & vision & vestibule Stand on pillow or spongy thing and close the eye .you are testing the vestibule alone you. If you did not find anything you can ask him to step. It is called "Vocada test".One of the vestibular examination

⁸ (has inner and outer hair cells → **responsible for hearing**) – (each part of the cochlea responds to specific kHz to conduct to the nerve) - responsible for linear acceleration


⁹ The saccule tells you when you stop moving and the utricle is responsible for head tilting) - linear acceleration

:Utricle & saccule and cristae of the Semicircular canals

?How many factors our balance depends on
 proprioception 2- vision 3- cerebellum -1



So, you have to make sure when someone ○
 came to you with imbalance it's not b/c of
 the cerebellum by testing it, then roll out the
 (peripheral (proprioception
 Testing the vision > by closing the eye -
 Testing the proprioception > by asking him -
 to stand on sponge When you do so (closing
 the pt. Eyes and making him stand on
 sponge) you're eliminating the vision and the
 proprioception effects and after it you can
 .make sure you're testing only the vestibule

In case of dizziness related to ear problem; it's either due to effect in the vestibular nerve (called -
 vestibular neuritis due to URI) or Benign positional vertigo (inside the vestibule there are fluid and
 gelatinous material that has Ca particles within it ; with minor trauma or any minor head concussion
 these Ca particles will go out from the gelatinous material to the fluid > once the pt. Moves his head up >
 movement of these Ca particles rapidly "when it was in the gelatinous material its movement was
 slowly" > vertigo not imbalance > treated by repositioning exercise after checking the type of nystagmus
 "horizontal = lateral Semicircular canal , rotatory= superior (geogravic) and posterior (ageogravic)
 (Semicircular canals the pt. have



:Cochlea

The cochlea is divided into 3 rooms (scala tympani, scala
 (media, scala vestibuli
 Most important is scala media; where hearing takes place. -
 .It contains hair cells and tectorial membrane
 The sounds wave vibrate the Tympanic membrane -> the -
 ossicles move (stapes act as a piston in scala vestibuli) it
 pushes the fluid away , and at the apical part (helicotrema)
 the fluid back to scala media so the wave of fluid will push
 the hair cells in it and it will touch tectorial membrane and
 will produce electrical stimulus and pass it to through the
 .spiral ganglion to the 8th nerve






Sound of **High frequency** affects the **basal portion of Cochlea**

Sounds of **Low frequency** affect the **apical portion of Cochlea**

B) Internal Auditory Canal


Contains: ○ Cochleovestibular nerve ○ Facial nerve




CENTRAL CONNECTIONS OF COCHLEAR NERVE	CENTRAL CONNECTIONS OF VESTIBULAR NERVE
<p>The principal human auditory cortex is located deep within the sylvian fissure on the superior surface of the temporal lobe</p> <p>The primary auditory cortex is often referred to as Brodmann .area 41</p> 	<p>PATHWAYS OF VESTIBULAR NERVE</p> 

➤ Blood supply of inner ear:

● Anterior inferior cerebellar artery → Labyrinthine artery → common cochlear and anterior vestibular




Physiology of the ear

	Function
External Ear	<ul style="list-style-type: none"> ● Protection of the middle ear: - Cerumen (wax), - Curvature. ● Auditory functions: – Sound Conduction. – Increase sound pressure by the resonance function
Eustachian Tube	<ul style="list-style-type: none"> ● Protection. ● Ventilation. ● Drainage 

<p>Middle Ear 11</p>	<ul style="list-style-type: none"> ● Conduction of sound¹⁰ ● Transformer mechanism: - Hydraulic action, - Ossicular leverage 1.3x amplification due to size difference between malleus and incus Stapes alone increase sound by 2000 hz, any problem in stapes we will find the audiogram fall at 2000 hz ● Protection to the inner ear. - Stapedial reflex, If the sound very loud it contracts to reduce the sound energy
<p>Inner Ear</p>	<ul style="list-style-type: none"> ● Hearing Function: ○ Transduction of sound to action potentials¹² Some cases they have the round window closed, once we open it they listen. ● Vestibular Function¹³ : <ul style="list-style-type: none"> ○ Participate in maintaining body balance, the mechanisms of maintaining body balance: (see up for more info) ○ Brain stem: is the center of balance. It's connected to: <ul style="list-style-type: none"> ■ Cerebellum to coordinate muscle tone and Cerebral cortex for the feeling of space. ■ Input: Proprioceptive (sensation), Visual and Vestibular ■ Output: gives information to: Postural muscles and Ocular muscle.



VESTIBULAR SYSTEM



¹⁰ there is amplification of the sounds when it transferred from big area to smaller one (tympanic membrane about 8mm and the stapes 2mm) also the difference in length b/w the malleus"longer" and incus"shorter" (about 1.3) increases the amplification; that's why the sound which we receive increases in almost 500-800 db - If there is reflection of the round window that's means the ossicle function is good,


¹¹ middle ear plays an important role in the process of impedance matching between the air-filled middle ear and the fluid-filled inner ear to allow for efficient sound transmission (**Impedance matching**-amplification-): Area ratio between the TM and the stapes footplate (20:1), Ossicular Coupling: lever ratio

¹² Any disease lead to closure of the round window will result in conductive hearing loss in the inner ear

¹³ Balance is central and peripheral. Central: brain stem, cerebellum and cerebral cortex. Peripheral: vision, vestibular and proprioception. How to examine it? Gait, Eye movement (nystagmus).

Development of the ear

- **External ear:** 1st pharyngeal cleft & arch
- **Middle ear:** 1st pharyngeal pouch & 1st and 2nd arches
- **Inner ear:** Ectoderm of hindbrain




Diseases of the external ear

congenital diseases

<p>:(Anotia (Atresia</p>	<p>It's the total absence of the auricle most often with ● .narrowing or absence of the external auditory meatus .(CT; to check if there is other malformations (internally ● Bone conduction is preserved Treatment: bone hearing aid .(B.A.H.A (on mastoid</p> <p>No auricles + canal atresia We reconstruct the ear (Prosthetic ear (usually tumor patients Good sensory hearing > bone hearing aid</p>	
<p>:Microtia</p>	<p>It's a condition in which the external portion of the ear ● (the auricle) is malformed. There is also narrowing or absence of the external auditory canal Any kind of remnant ●</p>	
<p>Accessory :auricle</p>	<p>.It's a type of ear anomaly in the tragus area ● Treatment: Plastic reconstruction, B.A.H.A ● (bone anchored hearing aid). It can present with no effect. usually in syndromes</p>	<p>Extra pic</p>
<p>Preauricular :sinus</p>	<p>It's a common congenital malformation characterized by a ● nodule, dent or dimple located anywhere adjacent to the .external ear Susceptible to infection ●</p> <ul style="list-style-type: none"> ● Management: systemic antibiotics. If an abscess is present, it must be incised and drained (If got infected twice you must take it out by first testing its pathway through methylene blue injection or CT scan with contrast, but in the time of inflammation we do incision and drainage.) Most common embryological defect Run in families Removed only if infected by removing the whole tract guided meth <p>dye.</p>	<p>Extra pic</p>

**Protruding Ear:
:Bat ear**


management: "cosmetic" Pinnaplasty or otoplasty. Do if after age of school. Note: There is no direct blood supply to the cartilage
 Antihelix pulls ear back while helix pushes it forward; ●
 Antihelix is absent
 By incisionless otoplasty (suture tie)



:Trauma to The Auricle ❖

- Lacerations
- Hematoma auris
 - Complication : cauliflower ear
- Treatment: Excise fibrous tissue - Apply pressure dressing - drain.


When we treat hematoma? Immediate incision and drainage! So, don't develop into **cauliflower ear (necrosed cartilage)**.



Cauliflower ear

❖ PERICHONDRIITIS OF THE PENNA:

- **Perichondritis** is inflammation of the perichondrium, a layer of connective tissue, which surrounds cartilage. (with spared lobule area)
- Usually follow trauma to the cartilage (hematoma auris, surgical "mastoid surgery", frostbite, burn) or otitis externa & piercing (particularly with the modern trend for multiple perforations that go through the cartilage).
- Commonly caused by **Pseudomonas**.
- Fever, pain, redness and swelling (causes narrowing and further low hearing level).
- **Treatment:** must be vigorous and immediately by parenteral antibiotics & Evacuation
 - (Any cartilaginous organ that forms a hematoma must be drained as early as possible)
 - If it is due to piercing the stud should be removed.



Cauliflower Ear

○ Complications of Perichondritis or Trauma:


- **Cauliflower ear (End stage of untreated haematoma)**. The ear can be exposed to trauma and lacerations leading to the **formation of Hematoma**, so if anything happens between the skin and cartilage → Hematoma (Number 1 killer of the cartilage, why?)



Because the blood will not be able to reach the cartilage) → Ischemia → Necrosis → Ear deformity

❖ **Otitis Externa:** An acute (Less than 3 months) or chronic (more than 3 months) infection of the whole or a part of the skin of the external ear canal. Any pathology affecting skin can also affect external ear.

- Organisms enter the apopilosebaceous unit by break in skin
- Commonly caused by fingernail or Q-tip to relieve itching
- Periosteal lining of bony canal displaced by swelling
- Subacute or chronic develops if AOE not treated adequately



○ **Pathophysiology:** – Aggressive washing of wax or retention water, Microtrauma (cotton swabs, fingernails).

Invective	Reactive
<ul style="list-style-type: none"> – Bacterial: Pseudomonas (commonly in immunocompromised like diabetic, post radio or chemotherapy and it has a very bad smell if it presents with Cholesteatoma), Staphylococcus aureus (furuncle) most common, like in swimming ear. Proteus mirabilis, Fungal: (newspaper appearance) Aspergillus – Niger (spores forming, hyphae), Candida albicans .(whitish and cheesy, cotton like) dx by seeing it Viral: Herpes Zoster¹⁴ ... Others – 	<ul style="list-style-type: none"> Seborrheic: A disease of the sebaceous glands ■ characterized by excessive secretion of sebum or an alteration in its quality, resulting in an oily coating, crusts, or scales on the skin. It's usually painless Eczematous/Dermatitis: A non-contagious ■ inflammation of the skin, characterized chiefly by .redness, itching

○ **Clinical features of Otitis Externa:**

- **Itching**
- **Pain:** could be very severe because of underlying cartilage, evoked by movement of the jaw, because the ear auricle and external canal is attached to the TMJ (temporomandibular joint) pain can radiate to the throat!
- **Fullness**
- **Tenderness and swelling**, absent in otitis media.
- **Otorrhea:** No discharge or very little and scanty, not mucoid. Large discharge in otitis media. (Not mucus discharge because the skin does not contain mucus-secreting cells. If the discharge doesn't contain mucus, then it is from the External ear however if it contains mucus it is originating from the middle ear)
- **Deafness:** deafness caused by external ear needs to be completely obstructed, which is rare in otitis externa.
- **Changes in the lumen and skin of EAM (external auditory meatus)**

Physical exam:

¹⁴ Ramsay Hunt syndrome (RHS): facial nerve palsy (weakness) and vesicles.

Redness, swelling, sometimes you can't see the TM because of the swelling, protrusion, discharge, preauricular or face or neck extension


Gently tug up and back: if true AOE, **patient will not tolerate**

Clean canal thoroughly and examine under Microscope

Clinical types of otitis externa

:(Localize O.E (furuncle ❖


.small rounded swelling in the external canal ○



Diffuse infective O.E.: swimming ear ❖

General narrowing of the canal. (on Ex we can't see the external canal b\c ○ of the edema) The canal will close, and you will not be able to pass anything through it

علشان كذا لما نبي ندخل لهم قطرات نستخدم شاش ندخله داخل الأذن ونصير نبلة بالقطرات علشان توصل داخل وقت الانفيكشن ما نسوي أي تدخل جراحي لأنه يسبب ادهيجن وفاييروسيس



❖ Otomycosis: fungal infection (More in those who take Abx for a long time)


- Fungal vs. Bacterial
- Fungal: Less pain, more itching & NO fever.

Management: suction then antifungal cream.



Black = Aspergillus Niger

White = Candida Albicans



:BULLOUS MYRINGITIS ❖

Inflammatory condition involves the lateral surface of the TM and the ○ .medial portion of the canal wall

It typically occurs in association with upper respiratory infections and is ○ .more common in winter

separation of one layer of the tympanic membrane "bullous" > viral)

(infection > pain

:Clinical manifestations ○

;Severe otalgia >

;Serosanguinous otorrhea >

.Hearing loss >

Treatment **includes analgesics, topical antibiotic/steroid drops** to ○

.prevent bacterial superinfection

Do not touch, if we open we will make it bacterial



The hallmark clinical finding is: **bull**a over the TM and medial canal with serous or serosanguinous fluid

Herpetic O.E: ◦ herpes zoster oticus is a specific form of herpes zoster ◈ that presents with pre-eruptive (“pre-herpetic”) lesion reactivated from either the trigeminal or cervical ganglions

characterized by: **PAINFUL vesicles** ◦

Management: Steroids + Acyclovir ◦

Complications: Facial n. paralyti ◦

Small vesicles + facial weakness = Ramsay Hunt syndrome or HSV

Eczematous and seborrheic: O.E. painless ◈


If the eczema is only in the canal, keep on mind tympanic membrane perforation due to discharge



Management (to all clinical types):

- History and Physical examination.
- antibiotics: Anti-pseudomonal drops – Ciprodex
- **Swab for culture and sensitivity for ABx.**
- **Ear toilet:** cleaning the ear. Meticulous debridement of debris, pus and cerumen
- **Keep the ear dry. Suction cleaning (especially the fungal infection = Suction, Suction, Suction)** the antifungal won't go inside so we have to take the debris out.

- **Ear wick**¹⁵ (best used after shower not in dry ear without pushing more than the length of the cotton > to avoid injury, infection and cotton dislodge¹⁶) This is an ear wick or sponge in the external ear canal for ear drops



Ear wick

- Local Medication and analgesia to control pain. Not all E.O need oral or parenteral tx.

- Systemic medications: as in diabetics.
- Surgery may be required in chronic cases and failure of treatment because there is usually thickening in of the skin and closure of the canal.

-Recommendations regarding prevention

- 1- Avoid instrumentation
- 2- Keep H₂O out of the ear when possible



- IN CASE OF:

- Aspergillus Niger → Give antifungal drops.



¹⁵ زي ما قلنا نستخدمه أحيانا لو كانت القناة ضيقة علشان ندخل قطرات الإذن، لو استخدمنا باكينق لازم نخليه أكثر من يومين ، ضروري نغيره 14 ولو ما غيرناه وكمل فيها يومين لازم نعطيه مضاد حيوي
¹⁶ و أهد سألك اقدر استخدم زيت الزيتون ، الإجابة بتكون نعم خصوصا لو كانت إذنههم الخارجية جافة جدا بس بشرط مايكون عندي ثقب لانه خطير، أيضا الخل يقدرن يستخدمونه بحالات معينة مثل تسوس الأذن بس بمقدار ١:١ ماء مع خل تقاح علشان يشيل كل السكين دبزي

■ Herpetic O.E Tx: → Acyclovir if < 3 days, Steroids to reduce inflammation.




❖ **Acute necrotizing (malignant is not a cancer) otitis externa** / Skull base Osteomyelitis (last approved name): **Important** Infection of the roof of the EAC and skull base affected

○ An acute **Pseudomonas** infection of the skin of the external ear canal (skull base), which spread to the adjacent bone. (Deep seated pain for more than a month).

-Life-threatening; osteomyelitis of temporal bone

-AOE can spread via fissures of Santorini or tympanomastoid fissure

Diagnosis : Ct scan to rule out other pathology such as cholesteatoma
PET scan to highlight the area



-Diabetes, advanced age, severe otalgia , granulation tissue ,cranial nerve involvement , radiology

○ It has a triad:

1- ear discharge "Several weeks of purulent otorrhea with granulations",

2- headache (esp at night),

3- Immunocompromised pt.

○ It occurs mostly in **elderly diabetic patients**. (Immunocompromised) Important!

Elderly with otorrhea and nocturnal headache and in examination granulation tissue and cranial palsy= necrotizing malignant otitis externa (skull base osteomyelitis)

○ **Severe otalgia**. Earache in early stage. > 1 month


○ Lower Cranial nerve palsies (VIII, IX, X, XI, XII) (check the gag reflex), and **25% VII**

○ No signs of acute inflammation & No swelling.

○ On Ex: Granulation tissue in EAC, sequestra and Foul-smelling discharge from the floor of the external Auditory canal.

○ It can infect the base of the skull, the cranium Causing meningitis, brain abscess.

Almost always caused by Pseudomonas; can be fungal in HIV –




○ Radiology: always we do CT although it doesn't tell us the definitive dx, that's why we rely on nuclear scan Bone¹⁷ (Petrous) scan to rule out osteomyelitis. Bony erosion on contrast-enhanced CT

- MRI useful for soft-tissue diagnosis, but not for F-U

- Bone scan is sensitive, but not specific (Tc-99m most sensitive)

○ Granulation tissue at the junction of the bony and cartilaginous portions of the canal + -immunocompromised pt → Dx as Malignant Otitis Externa!

- culture and biopsy



○ **Treatment:**

➤ **Control of diabetes** (most important part of treatment)

➤ **Anti-Pseudomonas antibiotics.** At least 6 weeks

➤ **Local treatment and debridement** anti-pseudomonal ear drops.

➤ **The role of surgery remains controversial** (mostly if there was a complication). Surgical treatment reserved for clear failures of above medical treatment

MISCELLANEOUS CONDITIONS OF THE EXTERNAL EAR:

¹⁷ Technetium 99: for diagnosis and gallium: for follow up. So we do them first both, the gallium we do it to know the baseline and after the treatment we do it "gallium" again to know the progress.

Wax:

- Mixture of ceruminous and sebaceous glands secretion

We always remove wax before examination






- Could be liquidy soft, scaly, hard
- Normally is expelled from the canal aided by movements of the jaw
- When accumulated it may cause deafness, earache or tinnitus
- Wax on tympanic membrane is very dangerous, it could be hiding retraction behind specially in parus flaccida or cholesteatoma

Treatment: is by removal using syringing very rare nowadays > anything you do it in ear will cause vasovagal + there will be stimulation to the lateral semicircular canal bc of the water temperature that we are using), suction or instrumentation

In irrigation we have to make sure of the temperature (same as body temperature) to avoid dizziness

crocodile forceps/ ear forceps

Hock



:KERATOSIS OBTURANS

○ Accumulation of desquamated epithelium (skin not wax) in the bony canal. (the difference b\w it and Cholesteatoma that in the later one we have normal skin in abnormal place)

It is excessive scaling of the skin causing very hard wax



Doesn't cause erosion only expansion (intact canal)

This is how we differentiate with external ear cholesteatoma

It may be associated with Sinusitis, Bronchiectasis, Primary ciliary dyskinesia. (it doesn't cause bony erosion but it lead to compression "pressure like effect" and widening of the canal)

○ Usually cause deafness and pain.

Treatment is periodic removal



18

Acute otitis media

Acute infection of the mucous membrane lining of the middle ear cleft.

● The definition is specific to infection because in chronic Otitis media it can be due to infection of normal inflammation.

● Predisposing factors:

○ **Age:** common in children as their Eustachian tube is more horizontal, wider and shorter in relation to their head.

○ **Male sex** ○ **Allergic Rhinitis** ○ **Smoking within the home**

○ **Bottle feeding:** more likely to have milk regurgitation (because children tend to drink while lying) in middle ear

○ **Climate** ○ **Crowded living conditions** (one infected will infect others) ○ **Heredity**

● Associated conditions:

cleft palate why? tensor palatini muscle is absent in cleft palate and its job to open ET when you swallow, immunodeficiency, ciliary dyskinesia, **Down syndrome**, and cystic fibrosis¹⁹.

Why in cleft palate? The muscles of the palate are affected and not well developed, so in cleft palate surgery ENT come to put tube to avoid otitis media with effusion for life

● **Route of infection:** ○ Eustachian tube. ○ External auditory canal(rupture): rare. ○ Blood borne.

● **Bacteriology:**

○ **Streptococcus pneumonia** (Most common)

¹⁸ Trick, External ear is very painful, middle ear is painless or less painful


¹⁹ fluids are thick in those pts it is like glue

- Haemophilus influenzae
- Branhamella (moraxella) catarrhalis
- Streptococcus pyogenes
- Staphylococcus aureus





First three are the main bacteria in upper airway, it is not wrong to put anywhere ear, nose throat infections

● Pathophysiology:

○ The patient has an antecedent event (viral URI or allergy) → the event results in Congestion of the respiratory mucosa of the nose, nasopharynx, and Eustachian tube → Congestion of the mucosa in the Eustachian tube obstructs the narrowest portion of the tube, the isthmus → obstruction of the isthmus causes negative pressure followed by accumulation of secretions produced by the mucosa of the middle ear → these secretions have no egress and accumulate in the middle ear space → viruses and bacteria that colonize the upper respiratory tract can reach the middle ear via aspiration, reflux, or insufflation → microbial growth in the middle ear secretions may result in suppuration.



● Clinical picture ●

<p>Tubal occlusion: produces early signs of acute otitis media. ○ Discomfort, autophony (feeling own sounds), retracted drum. (opposite of bulging) caused by pressure difference. There is mild deafness. Tinnitus in children, not adults ○ First thing to happen in otitis media is redness/congestion > bulge (severe pain) > rupture if untreated > pus > abnormal/normal healing or perforation</p>		
<p>Suppurative inflammation of the middle ear: Fever, severe earache, (deafness, congestion and bulging drum (pus behind it</p>	 <p style="text-align: right;">New pic</p>	
<p>Tympanic membrane rupture: Otorrhea, Temperature subside. & earache subside (pain relief), perforated drum and Mucopurulent (discharge (if not treated</p>	 <p style="text-align: right;">New pic</p>	
<p>Resolution: Either the rupture will persist, and it will discharge from time to time (chronic otitis media) Or close spontaneously ("retraction" (common Tympanosclerosis "if not treated will retract if it was severe > ○ adhesive otitis media (tympanic membrane reaching the promontory" or the cochlea</p>		

The patient can present to you at any stage (mostly the congestion and bulging) and the treatment will be the same. However, the complications are different ➤

The patient will be in severe pain before the rupture of tympanic membrane due to the nerve stimulation and irritation by tension ➤


:Complication of acute and chronic OM

Extracranial: Acute mastoiditis ,Chronic mastoiditis, Postauricular abscess ,Bezold abscess - Temporal abscess , Petrous apicitis ,Labyrinthine fistula ,**Facial nerve paralysis**

Acute suppurative labyrinthitis


Intracranial: Meningitis ,Brain abscess, Subdural empyema ,Epidural abscess -

Lateral sinus thrombosis , Otitic hydrocephalus,Encephalocele and cerebrospinal fluid leakage



:Treatment ●

- Symptomatic
- Antimicrobials.
 - Amoxicillin (1st line) if allergic to penicillin & cephalosporins you give clarithromycin
 - Amoxycillin/clavulanic acid (B-lactamase bacteria) 2nd line.
 - Trimethoprim-Sulfamethoxazole.
 - Cefaclor, cefixime.
 - Erythromycin-sulfisoxazole
- Decongestant. (opening in case of bulging with severe pain to relieve it and in congestion)
- Myringotomy +/- tube.
- Ear toilet and local antibiotics.
 - Bulging + severe pain + adult > open small opening to relieve the pain
 - If not > nasal steroidal spray so eustachian tube opens and remove the pus + oral ABx



● **Recurrent Acute Otitis Media:**

Three or more attacks over a 6-months period or (six attacks in a year).

- O.M. + diffusion > sterile fluid in the middle ear, or dysfunctioning Eustachian tube like in down syndrome or cleft palate.



● **Treatment:**

- Long-term low dose antimicrobials

-Ventilation tube insertion "it allows the air to enter the middle ear and drainage of fluid from the Eustachian tube" (Myringotomy with pressure equalization tube) Most common in acute otitis media after resolving there will be fluids.

Pediatrics last for 3-6 weeks if more it will affect speech, so we drain through eustachian tube (myringotomy) by putting a tube between the External Canal and middle ear


We put it in anterior inferior to avoid ossicles' injury



Extra pic

Note: Most OM in pediatrics is viral in origin, so we don't always need to rush to give Abx. Unless: high grade





Questions

1. Which structure is anterior to the external auditory canal?
 - A. Air
 - B. TMJ
 - C. Jugular bulb
 - D. Tympanic membrane

2. The outer layer of tympanic membrane is:
 - A. stratified squamous epithelium
 - B. ciliated columnar epithelium
 - C. cuboidal epithelium
 - D. non stratified squamous epithelium

3. Tonsillitis's referred earache is by which nerve?
 - A. Cervical II & III
 - B. V (Trigeminal) cranial nerve
 - C. IX (Glossopharyngeal) cranial nerve
 - D. X (vagus) cranial nerve

4. Which one of the following abnormalities is susceptible to infections?
 - A. Microtia
 - B. Accessory auricle
 - C. Preauricular sinus
 - D. Protruding ear

5. How would you treat Otomycosis?
 - A. Nothing, it's selflimiting
 - B. suction

- C. antibiotics
- D. suction the antifungal

6. Protruding ear is caused by the absence of which structure?

- A. helix
- B. tragus
- C. antihelix
- D. triangular fossa

Answers: 1:B 2:A 3:C 4:C 5:D 6:C