


The Nasal Septum

- Prof. Surayie H Al Dousary MD
- X Rhinology research Chair Director
 - X Head of ENT Department
 - X Head of Saudi ENT Society
- X Rhinology Fellowship Program Director
 - www.profseraye.com
 - @sdousary

The Nasal Septum Development



I. Cartilaginous Vault

II. Bony Vault




Cartilaginous Septum

- Septal (quadrilateral) cartilage
- The vomeronasal cartilages
- Medial crura of the alar cartilages




The Membranous Septum (Mobile Septum)

- Anterior to the end of the septal cartilage.
- It is formed by skin and subcutaneous tissue of the nasal columella.




Bony Septum

- Composed of two major elements:
 - The Vomer
 - The Perpendicular plate of the Ethmoid




The perpendicular Plate of the Ethmoid (Mesoethmoid)

- Ossification completed by 17th year of age.
- Replacement of cartilaginous septum with thin bone.
- At the nasal roof it articulates with the cribriform plate and extends as the crista galli



The Vomer

- Develops from **connective tissue** membrane on each side of the septal cartilage.
- The intervening **cartilage absorbed** completed by mid adult hood.




Inequality of Growth

- Creating **septal spur**.
- Septum Elevations and ridgelike **protuberances**



Asymmetry of the Nasal Septum

- Approximately **80 %** of humans have DNS
- Any parts of the septum except the **posterior border**
- A common area of deflection is along the **articulation** between the vomer and the perpendicular plate of the ethmoid
- DNS to one side or S shape to both side



Septal Deviations

- SSx: unilateral nasal obstruction, hyposmia, epistaxis, recurrent sinusitis
- Dx: anterior rhinoscopy, Endoscopy and CT PNS


Surgical Management

- **Submucous Resection(SMR):** obstructing cartilaginous and bony portion
- **Septoplasty:** removal of deviated cartilaginous and bony septum with **reinsertion** after remodeling and **repositioning** (preserves support system, less risk of perforation)

INDICATIONS

- Nasal obstruction (deviated nasal septum),
 - Epistaxis, chronic sinusitis (when septum is obstructing),
 - Access for transseptal sphenoidotomy,
 - Headache from an impacted spur
 - Septal neoplasia (rare)



Goals



EMERGENCIES NASAL OBSTRUCTION

Diagnosis	Emergency	Complications
Septal hematoma	Elevation of mucosal perichondrium with cartilage devascularization	Septal cartilage necrosis, development of a saddle-nose deformity
Septal abscess	Intracranial extension of infection	Septal cartilage necrosis, development of a saddle-nose deformity, cavernous sinus thrombosis, intracranial infection
Mucormycosis	Tissue destruction	Extension to brain or orbit


Septoplasty



FREER



COTTLE





Submucous Resection

Surgical correction of nasal valve deformities

- Widening the valve apex
 - Spreader grafts
 - Osteotomies
- Widening the valve angle
 - Flaring sutures
 - Suspension sutures
 - Butterfly grafts
- Stiffening the lateral crura
 - Alar batten grafts
 - Lateral crural J-flap

Area Approachable
by Submucous
Resection





Complications

- Psynchia
- Perforation,
- Saddle nose deformity (over resecting cartilage anteriorly),
- Cribriform plate fracture,
- Septal hematomas,
- Anosmia,
- Septal abscess,
- Bleeding

Psynechia

- Cause
- Manifestation
- Treatment



Septal Perforation

■ Cause

- Septoplasties (Most Common Cause, >50%),
- Infections
- (Tertiary Syphilis),
- Trauma (Nose Picking),
- Neoplasms,
- Granulomatous
- Disease,
- Vasculitis,
- Cocaine Abuse,
- Corticosteroid nasal spray

Diagnosis :

- Anterior rhinoscopy
- Biopsy of granulation tissue or abnormal mucosa



■ Manifestation

- Obstructive Sensation From Turbulent Flow, May Be Asymptomatic
- Crusting,
- Epistaxis,
- Whistling,

■ Treatment

1. Saline irrigation, emollients
2. Consider sliding or rotating mucoperichondrial flaps with or without a fascial graft; contraindicated for large perforations (approximately >2 cm of vertical height), cocaine abusers, malignancy, granulomatous or vascular diseases
3. Silastic Button



Sadel Nasal Deformity

- Cause
- Manifestation
- Treatment



Septal Hematoma



Symptoms and Signs

- Unilateral obstruction (may be bilateral),
- septal swelling
-




Complications

- Septal Abscess,
- Cavernous Sinus Thrombosis,
- Saddle Nose Deformity

Treatment

- Immediate evacuation of hematoma
- Nasal packing
- Antibiotic prophylaxis

FESS Land Marks (CLOSE)



Indications for ESS


- Chronic sinusitis,
- Complicated sinusitis,
- Recurrent acute sinusitis,
- Failed medical management of acute sinusitis,
- Fungal sinusitis
- Obstructive nasal polyposis
- Sinus mucoceles
- Remove foreign bodies
- Tumor excision,
- Transsphenoidal hypophysectomy
- Orbital decompression,
- Dacryocystorhinotomy,
- Orbital nerve decompression
- Grave's ophthalmopathy
- Choanal atresia repair
- CSF leak repair
- Control epistaxis
- Septoplasty,
- Turbinectomy

FESS Goals

- **Complete extirpation of all the disease**
- **Permanent drainage and ventilation of the affected sinuses**
- **Postoperative access to the previously diseased areas.**

Extended FESS

- CT Guided FESS
- Power Instrument
- Mini FESS



Polypectomy




Ethmoidectomy





- **Postoperative Care:**

- **Sinus Packing**
 - **Oral Antibiotics for a minimum of 2 weeks**
 - **Aggressive nasal hygiene to prevent adhesions (saline irrigations)**
 - **Nasal steroids**
 - **Nasal debridement at 1, 3, and 6 weeks**
- 



Excellent results

- 71% normal at one year
- Meta analysis 89% success
 - with 0.6% complications

FESS Orbital Complications

- **Blindness**

- Indirect injury (retrobulbar Hematoma)
- Direct injury to the optic nerve

- **Orbital Fat Penetration:**

increases risk of retrobulbar hematoma

Rx: recognize orbital fat (orbital fat floats);

avoid further trauma; may complete the FESS;

avoid tight nasal packing;



Observe for vision changes, proptosis, or restricted ocular gaze

- **Diplopia**: orbital muscle injury, most commonly from medial rectus and superior oblique muscles
- **Epiphora**: injury to lacrimal duct system, avoid operating anterior to the attachment of the uncinata;
Rx: observation initially, if no resolution then dacryocystorhinostomy

Retrobulbar Hematoma

- Pathophysiology: most commonly from retraction injury of the anterior ethmoid artery which causes increased orbital pressure that compresses the vascular supply to the optic nerve, also may occur from venous injury near the lamina papyracea
- Avoidance: maintain orientation and operate under direct vision, examine CT for dehiscence, correct coagulopathies

Turbinate Hypertrophy




Turbinate Hypertrophy

- **Causes**
 - Infection
 - Compensation
 - Dysfunctional
 - Allergies
- **Manifestation**
 - Nasal obstruction
 - Mouth Breathing
 - Cause manifestation




Turbinate Treatment

- Treat underlying cause
- Surgical treatment
 - SMR
 - Turbinoplasty
 - SMD
 - Somnoplasty RF
 - Turbenectomy
 - Ultrasonic reduction



TURBINATE REDUCTION GOALS

- Mucosal preservation
- Controlled reduction
- Submucous scarring to reduce the erectile nature of the mucosa
- Bony reduction when necessary
- Minimal complications



Preoperative Postoperative