

Infection Control

DR. MAZIN BARRY, MD, FRCPC, FACP, DTM&H

Infectious Disease Consultant

Assistant Professor of Medicine

Outline

- Hospital Acquired Infections (HAI)
- Hand Hygiene (HH)
- Isolation Precautions
- Others

Hospital Acquired Infections (HAI)

- At any time, over 1.4 million people worldwide are suffering from infections acquired in hospital
- Between 5% and 10% of patients admitted to hospitals acquire one or more infections
- **Causes more serious illness**
- Prolong hospital stay
- Long-term disability
- High additional financial burden
- High personal burden on patients and their families
- Deaths

Most frequent sites of infection and their risk factors

URINARY TRACT INFECTIONS

34%

Urinary catheter

Urinary invasive procedures

The age group that needs the catheter usually Advanced age
Severe underlying disease

Habitat for bacteria Urolithiasis

Urine stasis, may be asymptomatic and need routine screening Pregnancy
Diabetes

13%

LOWER RESPIRATORY TRACT INFECTIONS

Mechanical ventilation

Aspiration

Nasogastric tube

Central nervous system depressants

Antibiotics and anti-acids **and analgesia**

Prolonged health-care facilities stay

Malnutrition

Advanced age

Surgery

Immunodeficiency

Most common sites of health care-associated infection and the risk factors underlying the occurrence of infections

SURGICAL SITE INFECTIONS

Inadequate antibiotic prophylaxis

Incorrect surgical skin preparation

Inappropriate wound care

Surgical intervention duration

The deeper the wound the higher the risk Type of wound

Poor surgical asepsis

Immunosuppressive condition Diabetes

Nutritional state

Immunodeficiency

Lack of training and supervision

17%

14%

BLOOD INFECTIONS

Vascular catheter

Neonatal age

Critical care

Severe underlying disease

Neutropenia

Immunodeficiency **like cancer patient or HIV**

New invasive technologies

Lack of training and supervision

So prophylactic antibiotic should be used

Prevention of HAI


- Validated and standardized prevention strategies have been shown to reduce HAI
- At least 50% HAI could be prevented
- Most solutions are simple and not resource-demanding and can be implemented
- Hand hygiene, bundles **MOST IMPORTANT FACTOR!**

Colonization Versus Infection

- People who carry bacteria without evidence of infection (fever, increased WBC) are colonized
- If an infection develops, it is usually from bacteria that colonize patients
- Bacteria that colonize patients can be transmitted from one patient to another by hands of healthcare workers

Bacteria can be transmitted even if the patient is not infected. *The patient doesn't have any symptom*

The Iceberg Effect




The inanimate environment can facilitate transmission



Hand transmission

- Hands are the most common vehicle to transmit healthcare associated pathogens
- Transmission of microbiological organisms from one patient to another via healthcare worker's hands

The Carriers of Top Ten Infectious Diseases Germs




In US 20,000 cases of HAIs are directly related to poor hand hygiene annually.

Why should you clean your hands

- Any HCW involved in health care needs to be concerned about hand hygiene
- Other HCW hand hygiene concerns you as well
- You must perform hand hygiene to :
 - protect the patient against harmful microbes in your hands or present on your skin
 - protect yourself and the healthcare environment from harmful microbes

FIVE MOMENTS OF HAND HYGIENE

5 Moments of Hand hygiene




How to clean your hands

- Handrubbing with alcohol-based handrub is the preferred routine method of hand hygiene if hands are not visibly soiled
- Handwashing with soap and water – essential when hands are visibly dirty or visibly soiled (following exposure to body fluids)


How to handrub?

WITH ALCOHOL-BASED FORMULATION

1a



1b




Apply a palmful of the product in a cupped hand and cover all surfaces.

2



Rub hands palm to palm

3




right palm over left dorsum with interlaced fingers and vice versa

4




palm to palm with fingers interlaced

5



backs of fingers to opposing palms with fingers interlocked

6




rotational rubbing of left thumb clasped in right palm and vice versa

7




rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa

8




rinse hands with water

9



dry thoroughly with a single use towel

10




use towel to turn off faucet



20-30 sec

8




...once dry, your hands are safe.

How to handwash?


WITH SOAP AND WATER

0



Wet hands with water

1



apply enough soap to cover all hand surfaces.

Before all of this all jewelry and watches should be removed and he nails are cut and no nail polish

11



...and your hands are safe.



40-60 sec

Hand hygiene and glove use

- The use of gloves does not replace the need to clean the hands
- Remove gloves to perform Hand hygiene, when an indication occurs while wearing gloves
- Wear gloves only when indicated, otherwise they become a major risk for germ transmission

Should be removed before leaving the patient's room

ISOLATION PRECAUTIONS

Mode of transmission

- A microorganism may be spread by a single or multiple routes.
 - Contact, direct or indirect
 - Droplet
 - Airborne
 - Vector-borne (usually arthropod) and
 - Common environmental sources or vehicles - includes food-borne and waterborne, medications e.g., contaminated IV fluids

Types of Isolation Precautions

- Standard precautions **Hand hygien**
- Transmission-based precautions
 - Contact precautions
 - Airborne precautions
 - Droplet precautions

TRANSMISSION-BASED PRECAUTIONS -

Contact Precautions

- Infections spread by direct or indirect contact with patients or patient-care environment –C. difficile, MRSA, VRE, ESBL, CRE and MDR GNR
- Limit patient movement
- Private/SINGLE room or cohort with patients with same infection
- Wear disposable gown and gloves when entering the patient room
- Remove and discard used gown and gloves inside the patient room
- Wash hands immediately after leaving the patient room
- Use dedicated equipment if possible (e.g., stethoscope)

Contact precautions signs

On patient room door as a reminder



Visitors must report to Nursing Station before entering.



Perform hand hygiene before entering and before leaving room.



Wear gloves when entering room or cubicle, and when touching patient's intact skin, surfaces, or articles in close proximity



Wear gown when entering room or cubicle and whenever anticipating that clothing will touch patient items or potentially contaminated environmental surfaces.



Use patient-dedicated or single-use disposable shared equipment or clean and disinfect shared equipment (BP cuff, thermometers) between patients.

PRECAUCIONES DE CONTACTO

Los visitantes deben presentarse primero al puesto de enfermería antes de entrar. Lávese las manos. Póngase guantes al entrar al cuarto.

CONTACT PRECAUTIONS

To prevent the spread of infection,

ANYONE* ENTERING THIS ROOM MUST WEAR:

One use - disposable



Gloves ✓



Gown ✓

Applies whether or not contact with the patient or the patient's environment is anticipated.

*Patient visitors do not need to wear gloves and a gown, but must wash hands upon entering and leaving this room.

Questions? Please call the Department of Infection Control & Prevention at 936-0725




Droplet Precautions

- ✘ Reduce the risk of transmission by large particle droplets (larger than 5 μ in size).
- ✘ Requires close contact between the source person and the recipient
- ✘ Droplets usually travel 3 feet or less
- ✘ E.g., influenza, MERS-CoV, other respiratory viruses, rubella, parvovirus B19, mumps, *H. influenzae*, and *N. meningitidis*

Droplet Precautions cont.




- A private/single room or
- Cohort with patient with active infection with same microorganism
- Use a mask when entering the room especially within 3 feet of patient **Surgical mask**
- Limit movement and transport of the patient. Use a mask on the patient if they need to be moved and follow respiratory hygiene/cough etiquette

Droplet precautions signs



STOP **DROPLET PRECAUTIONS** **ALTO**


Visitors must report to Nursing Station before entering.

-  Perform hand hygiene before entering and before leaving room
-  Wear mask when entering room
Visitors and health care workers
-  Dietary may not enter
No debe entrar el dietista

PRECAUCIONES DE GOTAS DIMINUTAS

Los visitantes deben presentarse primero al puesto de enfermería antes de entrar. Lávese las manos. Póngase máscara al entrar al cuarto. No debe entrar el dietista.


HD 5225 Rev. 12/06




DROPLET PRECAUTIONS

To prevent the spread of infection,

ANYONE ENTERING THIS ROOM MUST WEAR:

 **Surgical Mask** ✓



N-95 Respirators should not be used for personal protection of patients in droplet precautions.

Questions? Please call the Department of Infection Control & Prevention at 936-0725  Vanderbilt Infection Control and Prevention

Airborne Precautions





- ✘ Tuberculosis, measles, varicella, MERS-CoV (severe)
- ✘ Place the patient in an airborne infection isolation room (AIIR)
Negative pressure room. AIIR = Airborne infection isolation room
- ✘ Pressure should be monitored with visible indicator **Outside the room**
- ✘ Use of respiratory protection (e.g., fit tested N95 respirator) or powered air-purifying respirator (PAPR) when entering the room
- ✘ Limit movement and transport of the patient. Use a mask on the patient if they need to be moved
And keep them last on the list for any radiological investigation or procedure
- ✘ Keep patient room door closed.

Airborne precautions signs



AIRBORNE INFECTION ISOLATION PRECAUTIONS

Visitors must report to Nursing Station before entering.


-  Perform hand hygiene before entering and before leaving room
-  Wear N95 respirator when entering room
Visitors see nurse for instruction on proper use.
-  Keep door closed
-  Dietary may not enter
No debe entrar el dietista

PRECAUCIONES AMBIENTALES

Los visitantes deben presentarse primero al puesto de enfermería antes de entrar. Lávese las manos. Póngase máscara N95 con filtro al entrar al cuarto. Mantenga la puerta cerrada. No debe entrar el dietista.

AIRBORNE PRECAUTIONS


To prevent the spread of infection,
**ANYONE* ENTERING THIS
ROOM MUST WEAR:**



N-95 Respirator ✓

Also ensure that the **door** to the patient's room **remains closed at all times.**

***Patient visitors should wear a blue surgical mask while in the patient's room.**

Questions? Please call the Department of Infection Control & Prevention at 936-0725  VanderbiltInfectionControl.com/Prevention

Safe injection practices

- Safe needle practice
- Reporting of needle stick and sharp injuries to infection control department

Serologies and Vaccination

- HBSAB titre (above 10) **Hepatitis B surface antibody titre**
- VZV **Varicella zooster virus**
- MMR **Measels - Mumps - Rubella**
- Td **Tittunes**
- Seasonal Influenza Vaccine
Should be given yearly due to changing viral strains